

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
8 March 2001 (08.03.2001)

PCT

(10) International Publication Number
WO 01/16346 A1

- (51) International Patent Classification⁷: C12P 7/22, (74) Agent: SEAY, Nicholas, J.; Quarles & Brady LLP, P.O. Box 2113, Madison, WI 53701-2113 (US).
C12N 9/02, 9/14, 1/20, 15/00
- (21) International Application Number: PCT/US00/23878
- (22) International Filing Date: 30 August 2000 (30.08.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 60/151,440 30 August 1999 (30.08.1999) US
- (71) Applicant (for all designated States except US): WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 Walnut Street, 13th Floor, P.O. Box 7365, Madison, WI 53707-7365 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): SUTHERS, Patrick, F. [US/US]; 806 Olin Avenue, Apartment 1, Madison, WI 53715 (US). CAMERON, Douglas, C. [US/US]; 3590 Rainier Lane, North Plymouth, MN 55447 (US).
- (81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PRODUCTION OF 3-HYDROXYPROPIONIC ACID IN RECOMBINANT ORGANISMS

(57) Abstract: The production of 3-hydroxypropionic acid (3-HP) from glycerol in a bacterial host is described. 3-HP is a useful feedstock for the production of polymeric materials. The genetic engineering of a bacterial host with two enzymes is sufficient to enable production of 3-HP. One enzyme is a glycerol dehydratase and the other is an aldehyde dehydrogenase.

WO 01/16346 A1

PRODUCTION OF 3-HYDROXYPROPIONIC ACID IN RECOMBINANT ORGANISMS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims priority from U.S. Provisional Patent Application S.N.

5 60/151,440 filed August 30, 1999.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

The research project which gave rise to the invention described in this patent application was supported by EPA grant R824726-01. The United States Government

10 may have certain rights in this invention.

BACKGROUND OF THE INVENTION

The technology of genetic engineering allows the transfer of genetic traits between species and permits, in particular, the transfer of enzymes from one species to others. These techniques have first reached commercialization in connection with high-value added products such as pharmaceuticals. The techniques of genetic engineering are equally applicable and cost effective when applied to genes and enzymes which can be used to make basic chemical feedstocks.

A metabolic pathway of interest exists in the bacteria *Klebsiella pneumoniae*, which has the ability to biologically produce 3 - hydroxypropionaldehyde from glycerol. Native microorganisms have the ability to produce 1,3 - propanediol from glycerol as well. Commercial interests are exploring the production of 1,3 - propanediol from glycerol or glucose, in recombinant organisms which have been engineered to express the enzymes necessary for 1,3 - propanediol production from other organisms.

3 - hydroxypropionic acid CAS registry Number [503-66-2] (abbreviated as 3-HP) is a three carbon non-chiral organic molecule. The IUPAC nomenclature name for

this molecule is propionic acid 3 - hydroxy. It is also known as 3 - hydroxypropionate, β - hydroxypropionic acid, β - hydroxypropionate, 3 - hydroxypropionic acid, 3 - hydroxypropanoate, hydracrylic acid, ethylene lactic acid, β -lactic acid and 2 - deoxyglyceric acid. Applications of 3-HP include the manufacture of absorbable
5 prosthetic devices and surgical sutures, incorporation into beta-lactams, production of acrylic acid, formation of trifluoromethylated alcohols or diols, polyhydroxyalkonates, and co-polymers with lactic acid. 3-HP for commercial use is now commonly produced by organic chemical syntheses. The 3-HP produced and sold by these methods is relatively expensive, and it would be cost prohibitive to use it for the production of
10 monomers for polymer production. As discussed below, some organisms are known to produce 3-HP. However, there is not yet available a catalog of genes from these organisms and thus the ability to synthesize 3-HP using the enzymes natively responsible for the synthesis of that molecule in the native hosts which produce it does not now exist.

15 In addition to its commercial utility, 3-HP it is found in a number of biological processes, notably including many naturally occurring bio-polymers. Poly(3 - hydroxybutyrate) (PHB) is the most abundant member of the microbial polyesters which contain hydroxy monomers termed polyhydroxyalkonates (PHAs). PHB has utility as a biodegradable thermoplastic material and the material was first produced industrially in
20 1982.

The majority of published research on PHA's that contain 3-HP has concentrated on two bacterial sources: *Ralstonia eutropha* ("*Alcaligenes eutrophus*") and *Pseudomonas oleovorans*. Both *Ralstonia eutropha* and *Pseudomonas oleovorans* are able to grow on a nitrogen free media containing 3 - hydroxy - propionic acid, 1,5 -
25 pentanediol or 1,7 - heptanediol. When 3-HP is the major hydroxy-acid added to the growth media, poly(3 - hydroxybutyrate - co - 3 - hydroxypropionic acid) is formed containing 7 mol % 3 - hydroxypropionic acid. These cells also store 3 mol %, 3 - hydroxypropionic acid poly(3 - butyrate - co - 3 - hydroxypropionic acid).

Recombinant systems have been used to create PHAs. An *E. coli* strain
30 engineered to express PHA synthase from either *Ralstonia eutropha* or *Zoogloea ramigera* produced poly(3 - hydroxypropionic acid) when feed 1,3 - propanediol.

- Skraly, F. A. "Polyhydroxyalkanoates Produced by Recombinant *E. coli*." Poster at Engineering Foundation Conference: Metabolic Engineering II, 1998. An *E. coli* strain that expressed PHA synthase (MBX820), when provided with the genes encoding glycerol dehydratase and 1,3 - propanediol dehydratase from *K. pneumonia*, and 4 - hydroxybutyral- CoA transferase from *Clostridium kluyveri*, synthesized PHB from glucose.

- Glycerol dehydratase, found in the bacterial pathway for the conversion of glycerol to 1,3 - propanediol, catalyzes the conversion of glycerol to 3 - hydroxypropionaldehyde and water. This enzyme has been found in a number of bacteria including strains of *Citrobacter*, *Klebsiella*, *Lactobacillus*, *Enterobacter* and *Clostridium*. In the 1,3 - propanediol pathway a second enzyme 1,3 - propanediol oxido-reductase (EC 1.1.202) reduces 3 - hydroxypropionaldehyde to 1,3 - propanediol in a NADH dependant reaction. The pathway for the conversion of glycerol to 1,3 - propanediol has been expressed in *E. coli*. Tong et al., Applied and Environmental Microbiology 57 (12) 3541-3546. The genes responsible for the production of 1,3 - propanediol were cloned from the *dha* regulon of *Klebsiella pneumoniae*. Glycerol is transported into the cell by the glycerol facilitator, and then converted into 3 - hydroxy - propionaldehyde by a coenzyme B₁₂- dependent dehydratase. *E. coli* lacks a native *dha* regulon, consequently *E. coli* cannot grow anaerobically on glycerol without an exogenous electron acceptor such as nitrate or fumarate.

- Aldehyde dehydrogenases are enzymes that catalyze the oxidation of aldehydes to carboxylic acids. The genes encoding non-specific aldehyde dehydrogenases have been identified in a wide variety of organisms e.g.; *ALDH2* from *Homo sapiens*, *ALD4* from *Saccharomyces cerevisiae*, and from *E. coli* both *aldA* and *aldB*, to name a few. These enzymes are classified by co-factor usage, most require either AND⁺, or NADP⁺ and some will use either co-factor. The genes singled out for mention here are able to act on a number of different aldehydes and it likely that they may be able to oxidize 3 - hydroxy - propionaldehyde to 3 - hydroxypropionic acid.

BRIEF SUMMARY OF THE INVENTION

The present invention is intended to permit the creation of a recombinant microbial host which is capable of synthesizing 3-HP from a starting material of glycerol or glucose. The glycerol or glucose is converted to 3 - hydroxypropionaldehyde (abbreviated as 3-HPA) which is then converted to 3-HP. This process requires the so-called *dhaB* gene from *Klebsiella pneumoniae* which encodes the enzyme glycerol dehydratase any one of four different aldehyde dehydrogenase genes to convert 3-HPA to 3-HP. The four aldehyde dehydrogenase genes used were *aldA* from the bacterium *E. coli*, *ALDH2* from humans, *ALD4* from the yeast *Saccharomyces cerevisiae*, and *aldB* from *E. coli*. The yeast gene appeared to give the best results.

It is an object of the present invention to provide a genetic construct which encodes glycerol dehydratase and aldehyde dehydrogenase enzymes necessary for the production of 3 - hydroxypropionic acid from glycerol.

It is also an object of the present invention to provide a method for the production of 3 - hydroxypropionic acid from glycerol.

Other features and advantages of the invention will be apparent from the following description of the preferred embodiment thereof and from the claims.

BRIEF DESCRIPTION OF THE SEVERAL VIEWS OF THE DRAWINGS

Not applicable.

DETAILED DESCRIPTION OF THE INVENTION

It is disclosed here that it is possible to introduce into a bacterial host genes encoding two enzymes and thus confer upon that host the ability to produce 3-HP from glycerol. The two necessary enzymes are glycerol dehydratase and aldehyde dehydrogenase. It is here reported that the two enzymes are both necessary and sufficient to enable a strain of a suitable host, such as a competent *E. coli* strain, to make 3-HP from glycerol. An exemplary gene encoding a glycerol dehydratase is known, the *dhaB* gene from *Klebsiella pneumoniae*, sequenced and rendered convenient to use. Several exemplary aldehyde dehydrogenases are known, and their sequences are

presented here. From this information, it becomes practical to confer upon a bacterial host the ability to convert glycerol into 3-HP in a commercially reasonable manner.

It was not apparent before the completion of the work described here that these two diverse enzymes could be produced in a common host to produce the ability to make 3-HP. There are many known aldehyde dehydrogenase enzymes and genes, and the enzymes are known to have varying substrate specificities and efficiencies. There was not evidence, prior to the work described here, that the aldehyde dehydrogenase enzyme would work on the 3-hydroxypropionaldehyde (3-HPA) substrate to create 3-HP. Without that knowledge, there was no data from which to predict the effectiveness of the 3-HP production studies described below. An additional uncertainty arises from the fact that the intermediate aldehyde, 3-HPA, is toxic to many bacterial host and thus the survival of the host is dependent upon the relative rates of enzymatic production and conversion of the aldehyde intermediate to non-toxic 3-HP.

A difficulty in the realization of the production of 3-HP desired here is that ribosome binding sites from non-native hosts are often ineffectual and lead to poor protein production and that many non-native promoters are often poorly transcribed and a bar to high protein expression. However, the inventors also recognized that a non-native promoter that is known to be very active and is inducible by the addition of a small molecule unrelated to the pathway being expressed is often a very efficient way to express and regulate the levels of enzymes expressed in hosts such as *E. coli*. To achieve high levels of regulated gene expression plasmids were constructed which placed the expression of all exogenous genes necessary for the production of 3 - hydroxypropionic acid from glycerol under the regulation of the *trc* promoter. The *trc* promoter, is efficient, not native to *E. coli*, and inducible by the addition of IPTG.

The present specification describes a genetic construct for use in the production of 3 - hydroxypropionic acid from glycerol. The genetic construct includes exemplary DNA sequences coding for the expression of a glycerol dehydratase and a DNA sequence coding for aldehyde dehydrogenase. The set of exemplary sequences necessary for the expression of glycerol dehydratase is collectively referred to as "*dhaB*". The set of sequences necessary for the expression of aldehyde dehydrogenase includes any one of four different genes which proved efficacious. The individual

aldehyde dehydrogenase sequences referred to individually as *ALDH4*, *ALD2*, *aldA* and *aldB*.

Producing 3 - hydroxypropionic acid in a foreign host

In the work described below, the enzymes necessary for the production of 3 - hydroxypropionic acid from glycerol in *E. coli* were expressed under the regulation of the *trc* promoter, a non-native promoter inducible by the addition of IPTG. The glycerol dehydratase was encoded by the *dhaB* gene from *Klebsiella pneumoniae*, the aldehyde dehydrogenases used was any one of four different genes (*ALDH2* from *Homo sapiens*, *ALD4* from *S. cerevisiae*, *aldB* from *E. coli* or *aldA* from *E. coli*). Expression of these genes coding for glycerol dehydratase and any one of the genes encoding an aldehyde dehydrogenases was sufficient to enable the construct to produce 3-HP when the fermentation media was supplemented with glycerol. In all of these constructs, the *dhaB* gene was downstream from the gene encoding the aldehyde dehydrogenase used, and expression of both genes was regulated by the *trc* promoter. This order, however, is not required and the order of the genes on a construct and the use of multiple constructs is possible.

In a minimal genetic construct made based on the data presented here, the only genetic elements present that would be necessary are the structural genes *dhaB* and an aldehyde dehydrogenase gene encoding a protein that efficiently catalyzes the oxidation of 3-hydroxypropionaldehyde to 3-hydroxypropionic acid, and non-native promoter sequences specifically selected to give the type of inducible control most appropriate for the context of the process in which the construct is to be used. Extraneous pieces of DNA, whether retained in the construct or added from other DNA sequences, would not necessarily be detrimental to effective 3-HP synthesis by the host organism, but would not be needed. Each sequence to be translated would necessarily be preceded by a ribosome binding site, functional in the selected host so that the messenger RNA(s) coding for the proteins of interest could be translated by ribosomes. Terminator sequences immediately downstream of each translated unit would also be necessary in some organisms, particularly in eukaryotes. The construct could be part of an autonomously replicating sequence, such as a plasmid or phage vector, or could be

integrated into the genome of the host.

The structural genes and appropriate promoter(s) could be isolated by the use of restriction enzymes, by the polymerase chain reaction (PCR), by chemical synthesis of the appropriate oligonucleotides, or by other methods apparent to those skilled in the art or molecular biology. The promoter(s) would be derived from genomic DNA of other organism or from artificial genetic constructs containing promoters. Appropriate promoter fragments would be ligated into the construct upstream of the structural genes in any one of several possible arrangements.

The aldehyde dehydrogenase expressed would have: high specific activity towards 3-hydroxypropionaldehyde; be very stable in the host it is expressed in; be readily over expressed in the selected host; not be inhibited by either the substrates necessary for the reaction or the products formed by the reaction; be fully active under the fermentation conditions most favorable for the production of 3 - hydroxypropionic acid and be able to use either NAD⁺ or NADP⁺.

One possible arrangement is the true operon, where one promoter is used to direct transcription in one direction of all necessary Open Reading Frames (ORFs). The entire message is then contained in one messenger RNA. The advantages of the operon are that it is relatively easy to construct, since only one promoter is needed; that it is relatively simple to replace the promoter with another promoter if that would be desirable later; and that it assures that the two genes are under the same regulation. The main disadvantage of the operon scheme is that the levels of the expression of the two genes cannot be varied independently. If it is found that the genes, for optimal 3 - hydroxypropionic acid synthesis, should be expressed at different levels, the operon in most cases cannot be used to realize this.

Another possible arrangement is the multiple-promoter scheme. Two or more promoters, with the same or distinct regulatory behavior, could be used to direct transcription of the genes. For example, one promoter could be used to direct transcription of *dhaB* and one to direct transcription of the gene encoding the appropriate aldehyde dehydrogenases. Because the genes theoretically can be transcribed and translated separately, a great number of combinations of multiple promoters is possible. Additionally, it would be most desirable to prevent the promoters

from interfering with one another. This could be achieved either by placing two promoters into the construct such that they direct transcription in opposite directions, or by inserting transcriptional terminator sequences downstream of each separately transcribed unit. The main advantage of the multiple-promoter construct is that it permits independent regulation of as many distinct units as desired, which could be important. The disadvantages are that it would be more difficult to construct; more difficult to amend later; and more difficult to effectively regulate, since multiple changes in fermentation conditions would need to be introduced and might render the performance of the fermentation somewhat less predictable.

- 10 In any construct, the promoter sequence(s) used should be functional in the selected host organism and preferably provide sufficient transcription of the genes comprising the glycerol to 3 - hydroxypropionic acid pathway to enable the construct to be adequately active in that host. The promoter sequence(s) used would also effect regulation of transcription of the genes enabling the glycerol to 3-HP pathway to be
- 15 adequately active under the fermentation conditions employed for 3-HP production, and preferably they would be inducible, such that expression of the genes could be modulated by the inclusion in, or exclusion from, the fermentation of a certain agents or conditions.

- A plausible example of the use of such a construct follows: one promoter, which
- 20 induced by the addition of an inexpensive chemical (the inducer) to the medium, could control transcription of both the *dhaB* gene and the gene encoding the appropriate aldehyde dehydrogenase. The cells would be permitted to grow in the absence of the inducer until they accumulated to a predetermined level. The inducer would then be added to the fermentation and nutritional changes commensurate with the altered
- 25 metabolism would be made to the medium as well. The cells would then be permitted to utilize the substrate(s) provided for 3-HP production (and additional biomass production if desired). After the cells could no longer use substrate to produce 3-HP, the fermentation would be stopped and the 3-HP recovered.

Genetic Sequences

- 30 To express glycerol dehydratase and a suitable aldehyde dehydrogenase, the two

enzymes necessary for the production of 3 - hydroxypropionic acid from glycerol, it is required that the DNA sequences containing the glycerol dehydratase and aldehyde dehydrogenase coding sequences be combined with at least a promoter sequence (preferably a non-native promoter although some native promoter activity may be present). An exemplary method of construction is described in the example below. To ensure that the present specification is enabling, the full sequences of the coding regions of genes for these enzymes is presented here.

Sequences 1, 3, 5 and 7 present different native genomic sequences for genes encoding aldehyde dehydrogenases.

10 SEQ ID NO:1 contains the full native DNA sequence encoding the *ALD4* enzyme from *Saccharomyces cerevisiae*. The amino acid sequence of the protein is presented as SEQ ID NO:2.

SEQ ID NO:3 includes the DNA sequence for the human *ALDH2* gene, again including the full protein coding region. The amino acid sequence for this human
15 alcohol dehydrogenase is presented in SEQ ID NO:4.

SEQ ID NO:5 and 7 respectively present the full coding sequences from the *E. coli* genes *aldA* and *aldB*, both of which encode alcohol dehydrogenases. The amino acid sequences for the proteins encoded by the genes are presented in SEQ ID NO: 6 and 8 respectively.

20 SEQ ID NO:9 contains the native genomic DNA sequence for the *dhaB* gene from the *dha* regulon of *Klebsiella pneumoniae*. The coding sequences for this complex regulon produces five polypeptides, which are presented as SEQ ID NOS:10 through 13, which together provide the activity of the glycerol dehydratase enzyme.

Each of these coding sequences can be used to make genetic constructs for the
25 expression of the appropriate enzymes in a heterologous hosts. In making genetic constructs for expression of the genes in such hosts, it is contemplated that heterologous promoters will be joined to the coding sequences for the enzymes, but all that it required is that the promoters be effective for the hosts in which the genes are to be expressed. It is also contemplated and envisioned that significant variations in DNA sequence are
30 possible from the native DNA coding sequences presented here. As is well known in the art, due to the degeneracy of the genetic code, many different DNA sequences can

encode the expression of the same protein. So, when this document uses language specifying a DNA sequence encoding a protein, it is intended to encompass any DNA sequence which can be used to express that protein even if different from the genomic sequences presented here. It is also contemplated that conservative changes in the amino acid sequences of the proteins specified here can be made without departing from the present invention. In particular, deletions, additions and substitutions of one or more amino acids in a protein sequence can almost always be made without changing protein functionality. When the name of a protein is used here, it is intended to be equally applicable to both such minor changes in amino acid sequence and to allelic variations in native protein sequence as occurs within the species named as well as other closely related species.

It is possible that many of the above DNA sequences could be truncated and still express a protein that has the same enzymatic properties. One skilled in the art of molecular biology would appreciate that minor deletions, additions and mutations may not change the attributes of the designated base pair sequences; many of the nucleotide of the designated base pair sequences are probably not essential for their unique function. To determine whether or not an altered sequence or sequences has sufficient homology with the designated base pairs to function identically, one would simply create the candidate mutation, deletion or alteration and create a gene construct including the altered sequence together with promoter and termination sequences. This gene construct could be tested as, described below, for the production of 3-HP from glycerol.

Certain DNA primers were used to isolate or clone the genomic DNA sequences used in the experiments described below. While the sequence information presented here is sufficient to enable the construction of expression plasmids incorporating the genes identified here, in order to redundantly enable the use of these genes, primers which may be used to isolate the genes from their native hosts are described below.

The primers aldA_L (SEQ ID NO:14), and aldA_R (SEQ ID NO:15), were used to amplify the 1513 bp *aldA* fragment from genomic *E. coli* DNA (strain MG1655, a gift from the Genetic Stock Center, New Haven, CT). The gel purified PCR fragment containing a DNA sequence coding for the expression of aldehyde dehydrogenase was

inserted into *NcoI-XhoI* site of pSE380 (Invitrogen, San Diego, CA) to give pPFS3. The resulting plasmid contained *aldA* under the control of the *trc* promoter. This construct allowed for high-level expression of the *aldA* gene from *E. coli* under regulation of the *trc* promoter. Unless indicated otherwise all molecular biology and plasmid constructions were done in *E. coli* AG1 (Stratagene, La Jolla, CA).

The primers aldB_L (SEQ ID NO:20) and aldB_R (SEQ ID NO:21), were used to amplify the 1574 bp *aldB* fragment from genomic *E. coli* DNA (strain MG1655). The resulting PCR converted the TGA stop codon into a TAA stop codon. The gel-purified PCR fragment containing the DNA sequence sufficiently coding for the expression of aldehyde dehydrogenase was inserted into the *KpnI-SacI* site of pSE380 to give pPFS12.

The primers ALD4_L (SEQ ID NO : 16), and ALD4_R (SEQ ID NO : 17), were used to amplify the 1595 bp ALD4 fragment from *S. cerevisiae* DNA (strain YPH500). The gel-purified fragment containing a DNA sequence coding for the expression of aldehyde dehydrogenase was inserted into the *KpnI-SacI* site of pPFS3 to give pPFS8. The resulting plasmid contained mature *ALD4* under control of the *trc* promoter.

The primers ALDH2_L (SEQ ID NO:18), and ALDH2_R (SEQ ID NO:19), were used to amplify the 1541 bp ALDH2 fragment from pT7-7::ALDH2, a gift from H. Weiner (Purdue University, West Lafayette, IN). The gel purified PCR fragment containing a DNA sequence sufficiently homologous to base pairs 22 to 1524, inclusive of SEQ ID NO : 3 so as to code for the expression of aldehyde dehydrogenase was inserted in to the *KpnI-SacI* site of pSE380 to give pPFS7. This sequence was moved from pPFS7 into the *KpnI-SacI* site of pPFS3 to give pPFS9. The resulting plasmid contained mature ALDH2 under the control of the *trc* promoter.

The primers pTRC_L (SEQ ID NO:22), and pTRC_R (SEQ ID NO:23), were used to amplify the 540 bp fragment from pSE380. The gel purified PCR fragment was inserted into the *HpaI-KpnI* site of pPFS3 to give pPFS13. The resulting plasmid deleted the "native" ribosome binding site of pSE380 and a *NcoI* site (which contained an extraneous ATG start codon upstream of the cloned genes). The *KpnI-SacI* fragments of pPFS8, pPFS9, and pPFS12 were inserted into the *KpnI-SacI* site of pPFS13 to give pPFS14, pPFS15, and pPFS16, respectively.

Assay for production of 3-HP

- The efficacy of changes made as contemplated herein can be checked by the following tests. To test for the production of 3-HP, fermentation products can be quantified with a Waters Alliance Integrity HPLC system (Milford, MA) equipped with
- 5 a refractive index detector, a photodiode array detector, and an Aminex HPX-87H (Bio-Rad, Hercules, CA) organic acids column. The mobile phase should be 0.01 N sulfuric acid solution (pH 2.0) at a flow rate of 0.5 mL/min. The column temperature should be set to 40°C. Compounds can be identified by determining if they co-elute with authentic standards. Prior to analysis, all samples should be filtered through 0.45 μ M
- 10 pore size membrane. (Gelman Sciences, Ann Arbor, MI). The fractions of the fermentation products collected using HPLC should be analyzed on a Varian Star 3400 CX, gas - chromatograph coupled to a Varian Saturn 3 mass spectrometer (GC-MS) (Walnut Creek, CA).

Assay for enzyme activity.

- 15 Aldehyde dehydrogenase activity can be determined by measuring the reduction of β -NAD⁺ at 25°C with 3 - hydroxypropionaldehyde as a substrate. All buffers should contain 1 mM ethylenediaminetetraacetic acid (EDTA), 0.1 mM Pefabloc SC (Boehringer Mannheim, Indianapolis, IN) and 1 mM Tris (carboxyethyl) phosphine hydrochloride (TCEP-HCL). For ALD4, the solution should contain 100 mM Tris HCL
- 20 Buffer (pH 8.0), 100 mM KCl. For ALDH2 the solution should contained 100 mM sodium pyrophosphate (pH 9.0). For AldA and AldB, the solution should contain 20 mM sodium glycine (pH 9.5). A total of 3.0 mL of buffer should be added to quartz cuvettes and allowed to equilibrate to assay temperature. From 5 to 20 μ L of cell extract should be added and background activity recorded after the addition of β -NAD⁺ to a
- 25 final concentration of 0.67 mM. The reaction should be started by the addition of substrate (either acetaldehyde, propionaldehyde, or 3 - hydroxypropionaldehyde) to a final concentration of 2 mM. Assay mixtures should be stirred with micro-stirrers during the assays.

For aldehyde dehydrogenase activity assays, one unit is defined as the reduction

- of 1.0 μ M of β -AND* per minute at 25° C. These reactions can be monitored by following the change in absorbance at 340 nm (A_{340}) at 25°C on a Varian Carry-1 Bio spectrophotometer (Sugar Land, TX). Total protein concentrations in the cell extracts can be determined using the Bradford assay method (Bio-Rad, Hercules, CA) with
- 5 bovine serum albumin as the standard.

EXAMPLES

Plasmid constructions.

- Klebsiella pneumoniae* expresses glycerol dehydratase, an enzyme that catalyzes the conversion of glycerol to 3 - hydroxypropionaldehyde, (*dhaB*) and 1,3 -
- 10 propanediol oxidoreductase an enzyme that catalyzes the conversion of 3 - hydroxypropionaldehyde to 1,3 - propanediol respectively (the gene product from *dhaT*). A plasmid encoding these two genes was created and expressed in *E. coli* (plasmid pTC53). The *dhaT* gene was deleted from pTC53 to create pMH34. The resulting plasmid still contained the DNA sequence complementary to base pairs 330 to
- 15 2153 inclusion of SEQ ID NO : 9, the complement of base pairs 2166 to 2591, inclusive, of SEQ ID NO : 9, and the complement of base pairs 3191 to 4858, inclusive, of SEQ ID NO : 9, so as to code for the expression of glycerol dehydratase. The fragment of DNA encoding these sequences was excised from pMH34 by cutting it with *SalI-XbaI*, and the resulting fragment was gel purified (the purified fragment was gift
- 20 from M. Hoffman of the University of Wisconsin - Madison). This DNA fragment was inserted into the *SalI-XbaI* site of pPFS13 to give pPFS17.

- The resulting plasmid contained both the *aldA* and *dhaB* genes under the control of the *trc* promoter. Similarly, the gel-purified *SalI-XbaI* fragment from pMH34 was inserted into the *SalI-XbaI* sites of pPFS14, pPFS15, and pPFS16 to give pPFS18,
- 25 pPFS19, and pPFS20, respectively. These plasmids contained *ALD4*, *ALDH2*, and *aldB*, respectively, as well as *dhaB* under the control of the *trc* promoter; in all of the constructs the *dhaB* gene were downstream of the gene encoding the aldehyde dehydrogenase.

Expression in *E. coli*

The efficacy of *E. coli* as a platform for the production of 3-HP from growth on glucose has been examined using a mathematical model developed for this purpose. The model was executed in two different ways assuming the conversion of one mole of glucose under either anaerobic or aerobic conditions either directly to 3-HP or to the production of 3-HP and ATP. The optimum yield under anaerobic conditions is 1 mole of 3-HP and 1 mole of lactate. The more realistic yield under anaerobic conditions is 0.5 moles of 3-HP, 1.5 moles of lactate and 1 mole of ATP. The optimum yield under aerobic conditions is 1.9 moles of 3-HP and 0.3 moles of CO₂. The realistic yield under aerobic conditions is 1.85 moles of 3-HP, 0.35 moles of CO₂ and 1 mole of ATP.

The effect of 3-HP concentration on *E. coli* strain MG1655 growth was measured. Cells were grown on standard media with and without the addition of up to 80g/L of 3-HP. The best fit of these data demonstrated that 3-HP was only 1.4 times as inhibitory as lactic acid on the growth of *E. coli*. It is possible to economically produce lactic acid using *E. coli*, since 3-HP is only 1.4 times more inhibitory than lactic acid, it should be possible to use *E. coli* as a host for the commercial production of 3-HP.

Media and growth conditions

The standard media contained the following per liter: 6 g Na₂HPO₄, 3 g KH₂PO₄, 1 g NH₄Cl, 0.5 g NaCl, 3 mg CaCl₂, 5 g yeast extract (Difco Laboratories, Detroit, MI) and 2 mM MgSO₄. When necessary to retain plasmids ampicillin (100 mg/mL) was added to the media. Isopropyl-β-thiogalactopyranoside (IPTG) was added in varying amounts to induce gene expression. All fermentations were carried out in an incubator-shaker at 37 C and 200 rpm. Anaerobic fermentations were carried out in 500-mL anaerobic flasks with 300 mL of working volume. Inocula for fermentations were grown overnight in Luria-Bertani medium supplemented with ampicillin is necessary. The 300-mL fermentations were inoculated with 1.5 mL of the overnight culture. For enzyme assays, fermentations were incubated for 24 hours.

Over expression of aldehyde dehydrogenase in *E. coli*

Cells were harvested by centrifugation at 3000 x g for 10 minutes at 4°C with a

Beckman (Fullerton, CA) model J2-21 centrifuge. Cell pellets were washed twice in 100 mM potassium phosphate buffer at pH 7.2 and re-suspended in appropriate assay resuspension buffer equal to 5 x of the volume of the wet cell mass. The cells were homogenized using a French pressure cell. The homogenate was centrifuged at 40000 x g for 30 minutes. The supernatant was dialyzed against the appropriate resuspension buffer using 10000 molecular weight cut-off pleated dialysis tubing (Pierce, Rockford, IL) at 4°C. Dialysis buffer was changed after 2 hours, and 4 hours, and dialysis was stopped after being allowed to proceed overnight.

E. coli AG1 cells transfected with the plasmids constructed to express the *aldA*, *ALD4*, *ALDH2*, or *aldB* genes were grown in 500-mL anaerobic flasks. Twelve hours after the fermentations were inoculated IPTG was added to induce enzyme expression. The cells were allowed to grow for an additional 12 hours then harvested and lysed as discussed above. The soluble fraction of the lysate was assayed for aldehyde dehydrogenase activity using the substrate 3-hydroxypropionaldehyde in the buffer appropriate for the particular enzyme expressed. The plasmid, aldehyde dehydrogenase expressed and specific activity measured (U/mg of protein) were as follows: pPFS13, *aldA*, 0.2; pPFS14, *ALD4*, 0.5, pPFS15, *ALDH2*, 0.3; and pPFS16, *aldB*, 0.1. The control, *E. coli* strain AG1 harboring plasmid pSE380, encoded no exogenous aldehyde dehydrogenase activity and it had no detectable activity with 3-HP as substrate. It is clear from the activity assays that all four aldehyde dehydrogenases were expressed in *E. coli*. The aldehyde dehydrogenase cloned from *Saccharomyces cerevisiae* (*ADH4*) had the highest activity when 3-hydroxypropionaldehyde was used as the substrate (0.5 units/mg of protein).

E. coli cells transformed with plasmids expressing: aldehyde dehydrogenase; both aldehyde dehydrogenase and glycerol dehydratase, or neither gene; were grown and assayed for their ability to produce 3-HP from glycerol. The cells were grown on standard media supplemented with 6 µM of Coenzyme B₁₂, under anaerobic conditions in the absence of light (to protect the integrity of the Coenzyme B₁₂ necessary for DhaB activity). After 12 hours, IPTG was added to induce expression of the genes under the *trc* promoter at the same time 5g/L of glycerol was added. After 12 more hours of anaerobic fermentation the fermentation broth was assayed for 3 - HP by HPLC and GC,

the plasmid, aldehyde dehydrogenase gene expressed and g/L of 3-HP measured were as follows: pSF17, *aldA*, 0.031; pPSF18 *ALD4*, 0.173; and pPSF19, *ALDH2*, 0.061. Cells expressing *dhaB* but no exogenous aldehyde dehydrogenase genes (plasmid pMH34) produced 0.015 g/L of 3-HP. Cells expressing *aldA*, *ALD4*, *ALDH2* or *aldB* 5 but not *dhaB* (plasmids pPFS13, pPFS14, pPFS15, pPFS16, respectively) all produced less than 0.005 g/L of 3-HP when the media the cells were growing in was supplemented with 2.5g/L of 3-hydroxypropionaldehyde.

Other Hosts and Promoters

Applications of the 3-hydroxypropionic acid pathway such as the genetic 10 constructs of the present invention can easily be expressed in other organisms. The required genes would need to be placed under control of an appropriate promoter or promoters. Some organism such as yeasts may require transcription terminators to be placed after each transcribed unit. The knowledge of the present invention makes such amendments possible. Such a genetic construct would need to be part of a vector that 15 could either replicate in the new host or integrate into the chromosome of the new host. Many such vectors are commercially available for expression in gram-negative and gram-positive bacteria, yeast, mammalian cells, insect cell, plant, etc. For example, to express the 3-hydroxypropionic acid pathway in *Rhodobacter capsulatus*, one could obtain vector pNH2 from the American Type Culture Collection (ATCC). This is a 20 shuttle vector for use in *R. capsulatus* and *E. coli*. Organisms such as *Saccharomyces cerevisiae* which can convert glucose to glycerol could be used as a host, such a construct would enable the production of 3-HP directly from glucose. Additionally, other substrates such as xylan could also be used given the selection of an appropriate host.

25 Stochiometric analysis shows that best stochiometric yield of 3-HP production in *E. coli* calculated on the basis of glucose consumed is obtained under aerobic conditions. Under aerobic condition CO₂ is the only carbon-containing co-product, in particular the generation of lactic acid which occurs under anaerobic conditions is avoided. Production of 3-HP under these conditions could result in a more economical 30 recovery of 3-HP from the fermentation broth.

Alternatively, the *dhaB* gene and a gene encoding the appropriate aldehyde dehydrogenase could be cloned into the multiple cloning site of this vector in *E. coli* to facilitate construction, and then transformed into *R. capsulatus*. The *R. capsulatus* *nifH* promoter, provided on the plasmid, could be used to direct the transcription in *R.*

- 5 *capsulatus* of the genes placed into pNF2 in series with one promoter, or with two copies of the *nifH* promoter. Expression of the genes in other organisms would require a procedure analogous to that presented here.

Alternative Aldehyde Dehydrogenases and Glycerol Dehydratases

- Applications of the pathway for the production of 3-hydroxypropionic acid from
10 glycerol can be made using other suitable aldehyde dehydrogenases. To be functional in this pathway an aldehyde dehydrogenase needs to be stable, readily expressed in the host of choice and have high enough activity towards 3-hydroxypropionaldehyde to enable it to make 3-HP. The knowledge of the present invention makes such amendments possible. A program of directed evolution could be undertaken to select
15 for suitable aldehyde dehydrogenases or they could be recovered from native sources, the genes encoding these enzymes in conjunction with a gene encoding an appropriate glycerol dehydratase activity, would then be made part of any of the constructs envisioned here to produce 3 - hydroxypropionic acid from glycerol.

- A similar program of enzyme improvement including for example directed
20 evolution could be carried out using the *dhaB* gene from *Klebsiella pneumoniae* as a starting point to obtain other variants of glycerol dehydratase that are superior in efficiency and stability to the form used in this invention. Alternatively, enzymes which catalyzes the same reaction may be isolated from others organisms and used in place of the *Klebsiella pneumoniae* glycerol dehydratase. Such enzymes may be especially
25 useful in alternative hosts wherein they may be more readily expressed, be more stable and more efficient under the fermentation conditions best suited to the growth of the construct and the production and recovery of 3-HP.

CLAIM OR CLAIMS

I/WE CLAIM:

1. A method for producing 3-hydroxypropionic acid comprising the steps of
providing in a fermenter a recombinant microorganism which expresses genes
5 for non-native enzymes which are capable of catalyzing the production of 3-
hydroxypropionic acid from glycerol;
providing a source of glycerol or glucose for the recombinant microorganism,
and
fermenting the microorganism under conditions which result in the accumulation
10 of 3-hydroxypropionic acid.
2. A method for producing 3-hydroxypropionic acid comprising the steps of
providing in a fermenter a recombinant microorganism which carries genetic
constructions for the expression of a glycerol dehydratase and an aldehyde
dehydrogenase which are capable of catalyzing the production of 3-hydroxypropionic
15 acid from glycerol;
providing a source of glycerol or glucose for the recombinant microorganism,
and
fermenting the microorganism under conditions which result in the accumulation
of 3-hydroxypropionic acid.

3. A method for producing 3-hydroxypropionic acid comprising the steps of providing in a fermenter a recombinant microorganism which carries a genetic construct which expresses the *dhaB* gene from *Klebsiella pneumoniae* and a gene for an aldehyde dehydrogenase, which are capable of catalyzing the production of 3-hydroxypropionic acid from glycerol;
- 5 providing a source of glycerol or glucose for the recombinant microorganism, and fermenting the microorganism under conditions which result in the accumulation of 3-hydroxypropionic acid.
- 10 4. The method of claim 3 wherein the gene for the aldehyde dehydrogenase is selected from the group consisting of *ALDH4*, *ALD2*, *aldA* and *aldB*.
5. The method of claim 3 wherein the aldehyde dehydrogenase is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.
- 15 6. A recombinant *E. coli* host comprising in its inheritable genetic materials foreign genes encoding a glycerol dehydratase and an aldehyde dehydrogenase, such that the host is capable of producing 3-hydroxypropionic acid from glycerol.
7. A recombinant *E. coli* host comprising in its inheritable genetic materials the *dhaB* gene from *Klebsiella pneumoniae* and the *ald4* gene from *Saccharomyces cerevisiae*, such that the host is capable of producing 3-hydroxypropionic from glycerol.

8. A bacterial host comprising in its inheritable genetic material a genetic construction encoding for the expression of a glycerol dehydratase enzyme and an aldehyde dehydrogenase enzyme, such that the bacterial host is capable of converting glycerol to 3-hydroxypropionic acid.

5 9. The bacterial host of claim 8 wherein the glycerol dehydratase from *Klebsiella pneumoniae*.

10 10. The bacterial host of claim 8 wherein the gene encoding the glycerol dehydratase is the *dhaB* gene from *Klebsiella pneumoniae*.

10 11. The bacterial host of claim 8 wherein the aldehyde dehydrogenase is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:4, SEQ ID NO:6 and SEQ ID NO:8.

12. The bacterial host of claim 8 wherein the gene for the aldehyde dehydrogenase is selected from the group consisting of *ALDH4*, *ALD2*, *aldA* and *aldB*.

SEQUENCE LISTING

<110> Suthers, Patrick F
Cameron, Douglas C.

<120> Production of 3-Hydroxypropionic Acid in Recombinant Organisms

5 <130> UW960296.96617

<140>

<141>

<160> 23

<170> PatentIn Ver. 2.1

10 <210> 1

<211> 1529

<212> DNA

<213> *Saccharomyces cerevisiae*

<220>

15 <221> CDS

<222> (25) .. (1509)

<400> 1

gtcgcggtac caaggaggta tcat atg tca cac ctt cct atg aca gtg cct 51

Met Ser His Leu Pro Met Thr Val Pro

20

1

5

atc aag ctg ccc aat ggg ttg gaa tat gag caa cca acg ggg ttg ttc 99

Ile Lys Leu Pro Asn Gly Leu Glu Tyr Glu Gln Pro Thr Gly Leu Phe

10

15

20

25

atc aac aac aag ttt gtt cct tct aaa cag aac aag acc ttc gaa gtc 147

25 Ile Asn Asn Lys Phe Val Pro Ser Lys Gln Asn Lys Thr Phe Glu Val

30

35

40

	att aac cct tcc acg gaa gaa gaa ata tgt cat att tat gaa ggt aga	195
	Ile Asn Pro Ser Thr Glu Glu Glu Ile Cys His Ile Tyr Glu Gly Arg	
	45 50 55	
	gag gac gat gtg gaa gag gcc gtg cag gcc gcc gac cgt gcc ttc tct	243
5	Glu Asp Asp Val Glu Glu Ala Val Gln Ala Ala Asp Arg Ala Phe Ser	
	60 65 70	
	aat ggg tct tgg aac ggt atc gac cct att gac agg ggt aag gct ttg	291
	Asn Gly Ser Trp Asn Gly Ile Asp Pro Ile Asp Arg Gly Lys Ala Leu	
	75 80 85	
10	tac agg tta gcc gaa tta att gaa cag gac aag gat gtc att gct tcc	339
	Tyr Arg Leu Ala Glu Leu Ile Glu Gln Asp Lys Asp Val Ile Ala Ser	
	90 95 100 105	
	atc gag act ttg gat aac ggt aaa gct atc tct tcc tgg aga gga gat	387
	Ile Glu Thr Leu Asp Asn Gly Lys Ala Ile Ser Ser Ser Arg Gly Asp	
15	110 115 120	
	gtt gat tta gtc atc aac tat ttg aaa tct tct gct ggc ttt gct gat	435
	Val Asp Leu Val Ile Asn Tyr Leu Lys Ser Ser Ala Gly Phe Ala Asp	
	125 130 135	
	aaa att gat ggt aga atg att gat act ggt aga acc cat ttt tct tac	483
20	Lys Ile Asp Gly Arg Met Ile Asp Thr Gly Arg Thr His Phe Ser Tyr	
	140 145 150	
	act aag aga cag cct ttg ggt gtt tgt ggg cag att att cct tgg aat	531
	Thr Lys Arg Gln Pro Leu Gly Val Cys Gly Gln Ile Ile Pro Trp Asn	
	155 160 165	
25	ttc cca ctg ttg atg tgg gcc tgg aag att gcc cct gct ttg gtc acc	579
	Phe Pro Leu Leu Met Trp Ala Trp Lys Ile Ala Pro Ala Leu Val Thr	
	170 175 180 185	

	ggt aac acc gtc gtg ttg aag act gcc gaa tcc acc cca ttg tcc gct	627
	Gly Asn Thr Val Val Leu Lys Thr Ala Glu Ser Thr Pro Leu Ser Ala	
	190 195 200	
	ttg tat gtg tct aaa tac atc cca cag gcg ggt att cca cct ggt gtg	675
5	Leu Tyr Val Ser Lys Tyr Ile Pro Gln Ala Gly Ile Pro Pro Gly Val	
	205 210 215	
	atc aac att gta tcc ggg ttt ggt aag att gtg gtt gag gcc att aca	723
	Ile Asn Ile Val Ser Gly Phe Gly Lys Ile Val Val Glu Ala Ile Thr	
	220 225 230	
10	aac cat cca aaa atc aaa aag gtt gcc ttc aca ggg tcc acg gct acg	771
	Asn His Pro Lys Ile Lys Lys Val Ala Phe Thr Gly Ser Thr Ala Thr	
	235 240 245	
	ggt aga cac att tac cag tcc gca gcc gca ggc ttg aaa aaa gtg act	819
	Gly Arg His Ile Tyr Gln Ser Ala Ala Ala Gly Leu Lys Lys Val Thr	
15	250 255 260 265	
	ttg gag ctg ggt ggt aaa tca cca aac att gtc ttc gcg gac gcc gag	867
	Leu Glu Leu Gly Gly Lys Ser Pro Asn Ile Val Phe Ala Asp Ala Glu	
	270 275 280	
	ttg aaa aaa gcc gtg caa aac att atc ctt ggt atc tac tac aat tct	915
20	Leu Lys Lys Ala Val Gln Asn Ile Ile Leu Gly Ile Tyr Tyr Asn Ser	
	285 290 295	
	ggt gag gtc tgt tgt gcg ggt tca agg gtg tat gtt gaa gaa tct att	963
	Gly Glu Val Cys Cys Ala Gly Ser Arg Val Tyr Val Glu Glu Ser Ile	
	300 305 310	
25	tac gac aaa ttc att gaa gag ttc aaa gcc gct tct gaa tcc atc aag	1011
	Tyr Asp Lys Phe Ile Glu Glu Phe Lys Ala Ala Ser Glu Ser Ile Lys	
	315 320 325	

	gtg ggc gac cca ttc gat gaa tct act ttc caa ggt gca caa acc tct	1059
	Val Gly Asp Pro Phe Asp Glu Ser Thr Phe Gln Gly Ala Gln Thr Ser	
	330 335 340 345	
	caa atg caa cta aac aaa atc ttg aaa tac gtt gac att ggt aag aat	1107
5	Gln Met Gln Leu Asn Lys Ile Leu Lys Tyr Val Asp Ile Gly Lys Asn	
	350 355 360	
	gaa ggt gct act ttg att acc ggt ggt gaa aga tta ggt agc aag ggt	1155
	Glu Gly Ala Thr Leu Ile Thr Gly Gly Glu Arg Leu Gly Ser Lys Gly	
	365 370 375	
10	tac ttc att aag cca act gtc ttt ggt gac gtt aag gaa gac atg aga	1203
	Tyr Phe Ile Lys Pro Thr Val Phe Gly Asp Val Lys Glu Asp Met Arg	
	380 385 390	
	att gtc aaa gag gaa atc ttt ggc cct gtt gtc act gta acc aaa ttc	1251
	Ile Val Lys Glu Glu Ile Phe Gly Pro Val Val Thr Val Thr Lys Phe	
15	395 400 405	
	aaa tct gcc gac gaa gtc att aac atg gcg aac gat tct gaa tac ggg	1299
	Lys Ser Ala Asp Glu Val Ile Asn Met Ala Asn Asp Ser Glu Tyr Gly	
	410 415 420 425	
	ttg gct gct ggt att cac acc tct aat att aat acc gcc tta aaa gtg	1347
20	Leu Ala Ala Gly Ile His Thr Ser Asn Ile Asn Thr Ala Leu Lys Val	
	430 435 440	
	gct gat aga gtt aat gcg ggt acg gtc tgg ata aac act tat aac gat	1395
	Ala Asp Arg Val Asn Ala Gly Thr Val Trp Ile Asn Thr Tyr Asn Asp	
	445 450 455	
25	ttc cac cac gca gtt cct ttc ggt ggg ttc aat gca tct ggt ttg ggc	1443
	Phe His His Ala Val Pro Phe Gly Gly Phe Asn Ala Ser Gly Leu Gly	
	460 465 470	

```

agg gaa atg tct gtt gat gct tta caa aac tac ttg caa gtt aaa gcg      1491
Arg Glu Met Ser Val Asp Ala Leu Gln Asn Tyr Leu Gln Val Lys Ala
      475                      480                      485

gtc cgt gcc aaa ttg gac gagtaagagc tcgaattcgc                        1529
5 Val Arg Ala Lys Leu Asp
      490                      495

<210> 2
<211> 495
<212> PRT
10 <213> Saccharomyces cerevisiae

<400> 2
Met Ser His Leu Pro Met Thr Val Pro Ile Lys Leu Pro Asn Gly Leu
      1                      5                      10                      15

Glu Tyr Glu Gln Pro Thr Gly Leu Phe Ile Asn Asn Lys Phe Val Pro
15                      20                      25                      30

Ser Lys Gln Asn Lys Thr Phe Glu Val Ile Asn Pro Ser Thr Glu Glu
      35                      40                      45

Glu Ile Cys His Ile Tyr Glu Gly Arg Glu Asp Asp Val Glu Glu Ala
      50                      55                      60

20 Val Gln Ala Ala Asp Arg Ala Phe Ser Asn Gly Ser Trp Asn Gly Ile
      65                      70                      75                      80

Asp Pro Ile Asp Arg Gly Lys Ala Leu Tyr Arg Leu Ala Glu Leu Ile
      85                      90                      95

Glu Gln Asp Lys Asp Val Ile Ala Ser Ile Glu Thr Leu Asp Asn Gly
25                      100                      105                      110

Lys Ala Ile Ser Ser Ser Arg Gly Asp Val Asp Leu Val Ile Asn Tyr
      115                      120                      125

```

Leu Lys Ser Ser Ala Gly Phe Ala Asp Lys Ile Asp Gly Arg Met Ile			
130	135	140	
Asp Thr Gly Arg Thr His Phe Ser Tyr Thr Lys Arg Gln Pro Leu Gly			
145	150	155	160
5 Val Cys Gly Gln Ile Ile Pro Trp Asn Phe Pro Leu Leu Met Trp Ala			
	165	170	175
Trp Lys Ile Ala Pro Ala Leu Val Thr Gly Asn Thr Val Val Leu Lys			
	180	185	190
Thr Ala Glu Ser Thr Pro Leu Ser Ala Leu Tyr Val Ser Lys Tyr Ile			
10	195	200	205
Pro Gln Ala Gly Ile Pro Pro Gly Val Ile Asn Ile Val Ser Gly Phe			
	210	215	220
Gly Lys Ile Val Val Glu Ala Ile Thr Asn His Pro Lys Ile Lys Lys			
225	230	235	240
15 Val Ala Phe Thr Gly Ser Thr Ala Thr Gly Arg His Ile Tyr Gln Ser			
	245	250	255
Ala Ala Ala Gly Leu Lys Lys Val Thr Leu Glu Leu Gly Gly Lys Ser			
	260	265	270
Pro Asn Ile Val Phe Ala Asp Ala Glu Leu Lys Lys Ala Val Gln Asn			
20	275	280	285
Ile Ile Leu Gly Ile Tyr Tyr Asn Ser Gly Glu Val Cys Cys Ala Gly			
	290	295	300
Ser Arg Val Tyr Val Glu Glu Ser Ile Tyr Asp Lys Phe Ile Glu Glu			
305	310	315	320
25 Phe Lys Ala Ala Ser Glu Ser Ile Lys Val Gly Asp Pro Phe Asp Glu			
	325	330	335

Ser Thr Phe Gln Gly Ala Gln Thr Ser Gln Met Gln Leu Asn Lys Ile
 340 345 350

 Leu Lys Tyr Val Asp Ile Gly Lys Asn Glu Gly Ala Thr Leu Ile Thr
 355 360 365

 5 Gly Gly Glu Arg Leu Gly Ser Lys Gly Tyr Phe Ile Lys Pro Thr Val
 370 375 380

 Phe Gly Asp Val Lys Glu Asp Met Arg Ile Val Lys Glu Glu Ile Phe
 385 390 395 400

 Gly Pro Val Val Thr Val Thr Lys Phe Lys Ser Ala Asp Glu Val Ile
 10 405 410 415

 Asn Met Ala Asn Asp Ser Glu Tyr Gly Leu Ala Ala Gly Ile His Thr
 420 425 430

 Ser Asn Ile Asn Thr Ala Leu Lys Val Ala Asp Arg Val Asn Ala Gly
 435 440 445

 15 Thr Val Trp Ile Asn Thr Tyr Asn Asp Phe His His Ala Val Pro Phe
 450 455 460

 Gly Gly Phe Asn Ala Ser Gly Leu Gly Arg Glu Met Ser Val Asp Ala
 465 470 475 480

 Leu Gln Asn Tyr Leu Gln Val Lys Ala Val Arg Ala Lys Leu Asp
 20 485 490 495

<210> 3

<211> 1541

<212> DNA

<213> Homo sapiens

<220>

<221> CDS

<222> (22)..(1521)

<400> 3

```

5 gcggtaccaa ggagatatca t atg tca gcc gcc gcc acc cag gcc gtg cct 51
      Met Ser Ala Ala Ala Thr Gln Ala Val Pro
      1             5             10

gcc ccc aac cag cag ccc gag gtc ttc tgc aac cag att ttc ata aac 99
Ala Pro Asn Gln Gln Pro Glu Val Phe Cys Asn Gln Ile Phe Ile Asn
10             15             20             25

aat gaa tgg cac gat gcc gtc agc agg aaa aca ttc ccc acc gtc aat 147
Asn Glu Trp His Asp Ala Val Ser Arg Lys Thr Phe Pro Thr Val Asn
      30             35             40

ccg tcc act gga gag gtc atc tgt cag gta gct gaa ggg gac aag gaa 195
15 Pro Ser Thr Gly Glu Val Ile Cys Gln Val Ala Glu Gly Asp Lys Glu
      45             50             55

gat gtg gac aag gca cgt gaa ggc cgc cgg ggc gcc ttc cag ctg ggc 243
Asp Val Asp Lys Ala Arg Glu Gly Arg Pro Gly Ala Phe Gln Leu Gly
      60             65             70

20 tca cct tgg cgc cgc atg gac gca tca cac agc ggc cgg ctg ctg aac 291
Ser Pro Trp Arg Arg Met Asp Ala Ser His Ser Gly Arg Leu Leu Asn
      75             80             85             90

cgc ctg gcc gat ctg atc gag cgg gac cgg acc tac ctg gcg gcc ttg 339
Arg Leu Ala Asp Leu Ile Glu Arg Asp Arg Thr Tyr Leu Ala Ala Leu
25             95             100             105

gag acc ctg gac aat ggc aag ccc tat gtc atc tcc tac ctg gtg gat 387
Glu Thr Leu Asp Asn Gly Lys Pro Tyr Val Ile Ser Tyr Leu Val Asp
      110             115             120

```

	ttg gac atg gtc ctc aaa tgt ctc cgg tat tat gcc ggc tgg gct gat	435
	Leu Asp Met Val Leu Lys Cys Leu Arg Tyr Tyr Ala Gly Trp Ala Asp	
	125 130 135	
	aag tac cac ggg aaa acc atc ccc att gac gga gac ttc ttc agc tac	483
5	Lys Tyr His Gly Lys Thr Ile Pro Ile Asp Gly Asp Phe Phe Ser Tyr	
	140 145 150	
	aca cgc cat gaa cct gtg ggg gtg tgc ggg cag atc att ccg tgg aat	531
	Thr Arg His Glu Pro Val Gly Val Cys Gly Gln Ile Ile Pro Trp Asn	
	155 160 165 170	
10	ttc ccg ctc ctg atg caa gca tgg aag ctg ggc cca gcc ttg gca act	579
	Phe Pro Leu Leu Met Gln Ala Trp Lys Leu Gly Pro Ala Leu Ala Thr	
	175 180 185	
	gga aac gtg gtt gtg atg aag gta gct gag cag aca ccc ctc acc gcc	627
	Gly Asn Val Val Val Met Lys Val Ala Glu Gln Thr Pro Leu Thr Ala	
15	190 195 200	
	ctc tat gtg gcc aac ctg atc aag gag gct ggc ttt ccc cct ggt gtg	675
	Leu Tyr Val Ala Asn Leu Ile Lys Glu Ala Gly Phe Pro Pro Gly Val	
	205 210 215	
	gtc aac att gtg cct gga ttt ggc ccc acg gct ggg gcc gcc att gcc	723
20	Val Asn Ile Val Pro Gly Phe Gly Pro Thr Ala Gly Ala Ala Ile Ala	
	220 225 230	
	tcc cat gag gat gtg gac aaa gtg gca ttc aca ggc tcc act gag att	771
	Ser His Glu Asp Val Asp Lys Val Ala Phe Thr Gly Ser Thr Glu Ile	
	235 240 245 250	
25	ggc cgc gta atc cag gtt gct gct ggg agc agc aac ctc aag aga gtg	819
	Gly Arg Val Ile Gln Val Ala Ala Gly Ser Ser Asn Leu Lys Arg Val	
	255 260 265	

	acc ttg gag ctg ggg ggg aag agc ccc aac atc atc atg tca gat gcc	867
	Thr Leu Glu Leu Gly Gly Lys Ser Pro Asn Ile Ile Met Ser Asp Ala	
	270 275 280	
	gat atg gat tgg gcc gtg gaa cag gcc cac ttc gcc ctg ttc ttc aac	915
5	Asp Met Asp Trp Ala Val Glu Gln Ala His Phe Ala Leu Phe Phe Asn	
	285 290 295	
	cag ggc cag tgc tgc tgt gcc ggc tcc cgg acc ttc gtg cag gag gac	963
	Gln Gly Gln Cys Cys Cys Ala Gly Ser Arg Thr Phe Val Gln Glu Asp	
	300 305 310	
10	atc tat gat gag ttt gtg gtg cgg agc gtt gcc cgg gcc aag tct cgg	1011
	Ile Tyr Asp Glu Phe Val Val Arg Ser Val Ala Arg Ala Lys Ser Arg	
	315 320 325 330	
	gtg gtc ggg aac ccc ttt gat agc aag acc gag cag ggg cgg cag gtg	1059
	Val Val Gly Asn Pro Phe Asp Ser Lys Thr Glu Gln Gly Pro Gln Val	
15	335 340 345	
	gat gaa act cag ttt aag aag atc ctc ggc tac atc aac acg ggg aag	1107
	Asp Glu Thr Gln Phe Lys Lys Ile Leu Gly Tyr Ile Asn Thr Gly Lys	
	350 355 360	
	caa gag ggg gcg aag ctg ctg tgt ggt ggg ggc att gct gct gac cgt	1155
20	Gln Glu Gly Ala Lys Leu Leu Cys Gly Gly Gly Ile Ala Ala Asp Arg	
	365 370 375	
	ggt tac ttc atc cag ccc act gtg ttt gga gat gtg cag gat ggc atg	1203
	Gly Tyr Phe Ile Gln Pro Thr Val Phe Gly Asp Val Gln Asp Gly Met	
	380 385 390	
25	acc atc gcc aag gag gag atc ttc ggg cca gtg atg cag atc ctg aag	1251
	Thr Ile Ala Lys Glu Glu Ile Phe Gly Pro Val Met Gln Ile Leu Lys	
	395 400 405 410	

```

    ttc aag acc ata gag gag gtt gtt ggg aga gcc aac aat tcc acg tac    1299
    Phe Lys Thr Ile Glu Glu Val Val Gly Arg Ala Asn Asn Ser Thr Tyr
                                415                420                425

    ggg ctg gcc gca gct gtc ttc aca aag gat ttg gac aag gcc aat tac    1347
    5 Gly Leu Ala Ala Ala Val Phe Thr Lys Asp Leu Asp Lys Ala Asn Tyr
                                430                435                440

    ctg tcc cag gcc ctc cag gcg ggc act gtg tgg gtc aac tgc tat gat    1395
    Leu Ser Gln Ala Leu Gln Ala Gly Thr Val Trp Val Asn Cys Tyr Asp
                                445                450                455

    10 gtg ttt gga gcc cag tca ccc ttt ggt ggc tac aag atg tgg ggg agt    1443
    Val Phe Gly Ala Gln Ser Pro Phe Gly Gly Tyr Lys Met Ser Gly Ser
                                460                465                470

    ggc cgg gag ttg ggc gag tac ggg ctg cag gca tac act gaa gtg aaa    1491
    Gly Arg Glu Leu Gly Glu Tyr Gly Leu Gln Ala Tyr Thr Glu Val Lys
    15 475                480                485                490

    act gtc aca gtc aaa gtg cct cag aag aac tcataagagc tcgaattcgc    1541
    Thr Val Thr Val Lys Val Pro Gln Lys Asn
                                495                500

    <210> 4
    20 <211> 500
    <212> PRT
    <213> Homo sapiens

    <400> 4
    Met Ser Ala Ala Ala Thr Gln Ala Val Pro Ala Pro Asn Gln Gln Pro
    25 1                5                10                15

    Glu Val Phe Cys Asn Gln Ile Phe Ile Asn Asn Glu Trp His Asp Ala
                                20                25                30

```


Val Ser Arg Lys Thr Phe Pro Thr Val Asn Pro Ser Thr Gly Glu Val			
35	40	45	
Ile Cys Gln Val Ala Glu Gly Asp Lys Glu Asp Val Asp Lys Ala Arg			
50	55	60	
5 Glu Gly Arg Pro Gly Ala Phe Gln Leu Gly Ser Pro Trp Arg Arg Met			
65	70	75	80
Asp Ala Ser His Ser Gly Arg Leu Leu Asn Arg Leu Ala Asp Leu Ile			
85	90	95	
Glu Arg Asp Arg Thr Tyr Leu Ala Ala Leu Glu Thr Leu Asp Asn Gly			
10 100	105	110	
Lys Pro Tyr Val Ile Ser Tyr Leu Val Asp Leu Asp Met Val Leu Lys			
115	120	125	
Cys Leu Arg Tyr Tyr Ala Gly Trp Ala Asp Lys Tyr His Gly Lys Thr			
130	135	140	
15 Ile Pro Ile Asp Gly Asp Phe Phe Ser Tyr Thr Arg His Glu Pro Val			
145 150	155	160	
Gly Val Cys Gly Gln Ile Ile Pro Trp Asn Phe Pro Leu Leu Met Gln			
165	170	175	
Ala Trp Lys Leu Gly Pro Ala Leu Ala Thr Gly Asn Val Val Val Met			
20 180	185	190	
Lys Val Ala Glu Gln Thr Pro Leu Thr Ala Leu Tyr Val Ala Asn Leu			
195	200	205	
Ile Lys Glu Ala Gly Phe Pro Pro Gly Val Val Asn Ile Val Pro Gly			
210	215	220	

Phe Gly Pro Thr Ala Gly Ala Ala Ile Ala Ser His Glu Asp Val Asp
 225 230 235 240

Lys Val Ala Phe Thr Gly Ser Thr Glu Ile Gly Arg Val Ile Gln Val
 245 250 255

5 Ala Ala Gly Ser Ser Asn Leu Lys Arg Val Thr Leu Glu Leu Gly Gly
 260 265 270

Lys Ser Pro Asn Ile Ile Met Ser Asp Ala Asp Met Asp Trp Ala Val
 275 280 285

Glu Gln Ala His Phe Ala Leu Phe Phe Asn Gln Gly Gln Cys Cys Cys
 10 290 295 300

Ala Gly Ser Arg Thr Phe Val Gln Glu Asp Ile Tyr Asp Glu Phe Val
 305 310 315 320

Val Arg Ser Val Ala Arg Ala Lys Ser Arg Val Val Gly Asn Pro Phe
 325 330 335

15 Asp Ser Lys Thr Glu Gln Gly Pro Gln Val Asp Glu Thr Gln Phe Lys
 340 345 350

Lys Ile Leu Gly Tyr Ile Asn Thr Gly Lys Gln Glu Gly Ala Lys Leu
 355 360 365

Leu Cys Gly Gly Gly Ile Ala Ala Asp Arg Gly Tyr Phe Ile Gln Pro
 20 370 375 380

Thr Val Phe Gly Asp Val Gln Asp Gly Met Thr Ile Ala Lys Glu Glu
 385 390 395 400

Ile Phe Gly Pro Val Met Gln Ile Leu Lys Phe Lys Thr Ile Glu Glu
 405 410 415

Val Val Gly Arg Ala Asn Asn Ser Thr Tyr Gly Leu Ala Ala Ala Val
 420 425 430

Phe Thr Lys Asp Leu Asp Lys Ala Asn Tyr Leu Ser Gln Ala Leu Gln
 435 440 445

5 Ala Gly Thr Val Trp Val Asn Cys Tyr Asp Val Phe Gly Ala Gln Ser
 450 455 460

Pro Phe Gly Gly Tyr Lys Met Ser Gly Ser Gly Arg Glu Leu Gly Glu
 465 470 475 480

Tyr Gly Leu Gln Ala Tyr Thr Glu Val Lys Thr Val Thr Val Lys Val
 10 485 490 495

Pro Gln Lys Asn
 500

<210> 5
 <211> 1512
 15 <212> DNA
 <213> Escherichia coli

<220>
 <221> CDS
 <222> (37)..(1473)

20 <400> 5
 gctaccatgg cttaaccggt accaaggaga tatcat atg tca gta ccc gtt caa 54
 Met Ser Val Pro Val Gln
 1 5

cat cct atg tat atc gat gga cag ttt gtt acc tgg cgt gga gac gca 102
 25 His Pro Met Tyr Ile Asp Gly Gln Phe Val Thr Trp Arg Gly Asp Ala
 10 15 20

```

    tgg att gat gtg gta aac cct gct aca gag gct gtc att tcc cgc ata   150
    Trp Ile Asp Val Val Asn Pro Ala Thr Glu Ala Val Ile Ser Arg Ile
          25                      30                      35

    ccc gat ggt cag gcc gag gat gcc cgt aag gca atc gat gca gca gaa   198
5  Pro Asp Gly Gln Ala Glu Asp Ala Arg Lys Ala Ile Asp Ala Ala Glu
      40                      45                      50

    cgt gca caa cca gaa tgg gaa gcg ttg cct gct att gaa cgc gcc agt   246
    Arg Ala Gln Pro Glu Trp Glu Ala Leu Pro Ala Ile Glu Arg Ala Ser
      55                      60                      65                      70

10  tgg ttg cgc aaa atc tcc gcc ggg atc cgc gaa cgc gcc agt gaa atc   294
    Trp Leu Arg Lys Ile Ser Ala Gly Ile Arg Glu Arg Ala Ser Glu Ile
          75                      80                      85

    agt gcg ctg att gtt gaa gaa ggg ggc aag atc cag cag ctg gct gaa   342
    Ser Ala Leu Ile Val Glu Glu Gly Gly Lys Ile Gln Gln Leu Ala Glu
15          90                      95                      100

    gtc gaa gtg gct ttt act gcc gac tat atc gat tac atg gcg gag tgg   390
    Val Glu Val Ala Phe Thr Ala Asp Tyr Ile Asp Tyr Met Ala Glu Trp
          105                      110                      115

    gca cgg cgt tac gag ggc gag att att caa agc gat cgt cca gga gaa   438
20  Ala Arg Arg Tyr Glu Gly Glu Ile Ile Gln Ser Asp Arg Pro Gly Glu
      120                      125                      130

    aat att ctt ttg ttt aaa cgt gcg ctt ggt gtg act acc ggc att ctg   486
    Asn Ile Leu Leu Phe Lys Arg Ala Leu Gly Val Thr Thr Gly Ile Leu
      135                      140                      145                      150

25  cgg tgg aac ttc ccg ttc ttc ctc att gcc cgc aaa atg gct ccc gct   534
    Pro Trp Asn Phe Pro Phe Phe Leu Ile Ala Arg Lys Met Ala Pro Ala
          155                      160                      165

```

	ctt ttg acc ggt aat acc atc gtc att aaa cct agt gaa ttt acg aca	582
	Leu Leu Thr Gly Asn Thr Ile Val Ile Lys Pro Ser Glu Phe Thr Thr	
	170 175 180	
	aac aat gcg att gca ttc gcc aaa atc gtc gat gaa ata ggc ctt cgg	630
5	Asn Asn Ala Ile Ala Phe Ala Lys Ile Val Asp Glu Ile Gly Leu Pro	
	185 190 195	
	cgc gcc gtg ttt aac ctt gta ctg ggg cgt ggt gaa acc gtt ggg caa	678
	Arg Gly Val Phe Asn Leu Val Leu Gly Arg Gly Glu Thr Val Gly Gln	
	200 205 210	
10	gaa ctg gcg ggt aac cca aag gtc gca atg gtc agt atg aca ggc agc	726
	Glu Leu Ala Gly Asn Pro Lys Val Ala Met Val Ser Met Thr Gly Ser	
	215 220 225 230	
	gtc tct gca ggt gag aag atc atg gcg act gcg gcg aaa aac atc acc	774
	Val Ser Ala Gly Glu Lys Ile Met Ala Thr Ala Ala Lys Asn Ile Thr	
15	235 240 245	
	aaa gtg tgt ctg gaa ttg ggg ggt aaa gca cca gct atc gta atg gac	822
	Lys Val Cys Leu Glu Leu Gly Gly Lys Ala Pro Ala Ile Val Met Asp	
	250 255 260	
	gat gcc gat ctt gaa ctg gca gtc aaa gcc atc gtt gat tca cgc gtc	870
20	Asp Ala Asp Leu Glu Leu Ala Val Lys Ala Ile Val Asp Ser Arg Val	
	265 270 275	
	att aat agt ggg caa gtg tgt aac tgt gca gaa cgt gtt tat gta cag	918
	Ile Asn Ser Gly Gln Val Cys Asn Cys Ala Glu Arg Val Tyr Val Gln	
	280 285 290	
25	aaa ggc att tat gat cag ttc gtc aat cgg ctg ggt gaa gcg atg cag	966
	Lys Gly Ile Tyr Asp Gln Phe Val Asn Arg Leu Gly Glu Ala Met Gln	
	295 300 305 310	

	gcg gtt caa ttt ggt aac ccc gct gaa cgc aac gac att gcg atg ggg	1014
	Ala Val Gln Phe Gly Asn Pro Ala Glu Arg Asn Asp Ile Ala Met Gly	
	315 320 325	
	ccg ttg att aac gcc gcg gcg ctg gaa agg gtc gag caa aaa gtg gcg	1062
5	Pro Leu Ile Asn Ala Ala Ala Leu Glu Arg Val Glu Gln Lys Val Ala	
	330 335 340	
	cgc gca gta gaa gaa ggg gcg aga gtg gcg ttc ggt ggc aaa gcg gta	1110
	Arg Ala Val Glu Glu Gly Ala Arg Val Ala Phe Gly Gly Lys Ala Val	
	345 350 355	
10	gag ggg aaa gga tat tat tat ccg ccg aca ttg ctg ctg gat gtt cgc	1158
	Glu Gly Lys Gly Tyr Tyr Tyr Pro Pro Thr Leu Leu Leu Asp Val Arg	
	360 365 370	
	cag gaa atg tcg att atg cat gag gaa acc ttt ggc ccg gtg ctg cca	1206
	Gln Glu Met Ser Ile Met His Glu Glu Thr Phe Gly Pro Val Leu Pro	
15	375 380 385 390	
	gtt gtc gca ttt gac acg ctg gaa gat gct atc tca atg gct aat gac	1254
	Val Val Ala Phe Asp Thr Leu Glu Asp Ala Ile Ser Met Ala Asn Asp	
	395 400 405	
	agt gat tac ggc ctg acc tca tca atc tat acc caa aat ctg aac gtc	1302
20	Ser Asp Tyr Gly Leu Thr Ser Ser Ile Tyr Thr Gln Asn Leu Asn Val	
	410 415 420	
	gcg atg aaa gcc att aaa ggg ctg aag ttt ggt gaa act tac atc aac	1350
	Ala Met Lys Ala Ile Lys Gly Leu Lys Phe Gly Glu Thr Tyr Ile Asn	
	425 430 435	
25	cgt gaa aac ttc gaa gct atg caa ggc ttc cac gcc gga tgg cgt aaa	1398
	Arg Glu Asn Phe Glu Ala Met Gln Gly Phe His Ala Gly Trp Arg Lys	
	440 445 450	

```

tcc ggt att ggc ggc gca gat ggt aaa cat ggc ttg cat gga tat ctg   1446
Ser Gly Ile Gly Gly Ala Asp Gly Lys His Gly Leu His Gly Tyr Leu
455               460               465               470

cag acc cag gtg gtt tat tta cag tct taagagctcg aattcccgtc   1493
5 Gln Thr Gln Val Val Tyr Leu Gln Ser
      475

gaaggctcta gactcgagcg   1513

<210> 6
<211> 479
10 <212> PRT
    <213> Escherichia coli

<400> 6
Met Ser Val Pro Val Gln His Pro Met Tyr Ile Asp Gly Gln Phe Val
  1               5               10               15

15 Thr Trp Arg Gly Asp Ala Trp Ile Asp Val Val Asn Pro Ala Thr Glu
      20               25               30

Ala Val Ile Ser Arg Ile Pro Asp Gly Gln Ala Glu Asp Ala Arg Lys
      35               40               45

Ala Ile Asp Ala Ala Glu Arg Ala Gln Pro Glu Trp Glu Ala Leu Pro
20      50               55               60

Ala Ile Glu Arg Ala Ser Trp Leu Arg Lys Ile Ser Ala Gly Ile Arg
      65               70               75               80

Glu Arg Ala Ser Glu Ile Ser Ala Leu Ile Val Glu Glu Gly Gly Lys
      85               90               95

25 Ile Gln Gln Leu Ala Glu Val Glu Val Ala Phe Thr Ala Asp Tyr Ile
      100               105               110

```

Asp Tyr Met Ala Glu Trp Ala Arg Arg Tyr Glu Gly Glu Ile Ile Gln
 115 120 125

Ser Asp Arg Pro Gly Glu Asn Ile Leu Leu Phe Lys Arg Ala Leu Gly
 130 135 140

5 Val Thr Thr Gly Ile Leu Pro Trp Asn Phe Pro Phe Phe Leu Ile Ala
 145 150 155 160

Arg Lys Met Ala Pro Ala Leu Leu Thr Gly Asn Thr Ile Val Ile Lys
 165 170 175

Pro Ser Glu Phe Thr Thr Asn Asn Ala Ile Ala Phe Ala Lys Ile Val
 10 180 185 190

Asp Glu Ile Gly Leu Pro Arg Gly Val Phe Asn Leu Val Leu Gly Arg
 195 200 205

Gly Glu Thr Val Gly Gln Glu Leu Ala Gly Asn Pro Lys Val Ala Met
 210 215 220

15 Val Ser Met Thr Gly Ser Val Ser Ala Gly Glu Lys Ile Met Ala Thr
 225 230 235 240

Ala Ala Lys Asn Ile Thr Lys Val Cys Leu Glu Leu Gly Gly Lys Ala
 245 250 255

Pro Ala Ile Val Met Asp Asp Ala Asp Leu Glu Leu Ala Val Lys Ala
 20 260 265 270

Ile Val Asp Ser Arg Val Ile Asn Ser Gly Gln Val Cys Asn Cys Ala
 275 280 285

Glu Arg Val Tyr Val Gln Lys Gly Ile Tyr Asp Gln Phe Val Asn Arg
 290 295 300

Leu Gly Glu Ala Met Gln Ala Val Gln Phe Gly Asn Pro Ala Glu Arg
 305 310 315 320

Asn Asp Ile Ala Met Gly Pro Leu Ile Asn Ala Ala Ala Leu Glu Arg
 325 330 335

5 Val Glu Gln Lys Val Ala Arg Ala Val Glu Glu Gly Ala Arg Val Ala
 340 345 350

Phe Gly Gly Lys Ala Val Glu Gly Lys Gly Tyr Tyr Tyr Pro Pro Thr
 355 360 365

Leu Leu Leu Asp Val Arg Gln Glu Met Ser Ile Met His Glu Glu Thr
 10 370 375 380

Phe Gly Pro Val Leu Pro Val Val Ala Phe Asp Thr Leu Glu Asp Ala
 385 390 395 400

Ile Ser Met Ala Asn Asp Ser Asp Tyr Gly Leu Thr Ser Ser Ile Tyr
 405 410 415

15 Thr Gln Asn Leu Asn Val Ala Met Lys Ala Ile Lys Gly Leu Lys Phe
 420 425 430

Gly Glu Thr Tyr Ile Asn Arg Glu Asn Phe Glu Ala Met Gln Gly Phe
 435 440 445

His Ala Gly Trp Arg Lys Ser Gly Ile Gly Gly Ala Asp Gly Lys His
 20 450 455 460

Gly Leu His Gly Tyr Leu Gln Thr Gln Val Val Tyr Leu Gln Ser
 465 470 475

<210> 7
 <211> 1574
 <212> DNA
 <213> *Escherichia coli*

5 <220>
 <221> CDS
 <222> (22)..(1557)

<400> 7
 gcggtaccaa ggaggatca t atg acc aat aat ccc cct tca gca cag att 51
 10 Met Thr Asn Asn Pro Pro Ser Ala Gln Ile
 1 5 10

aag ccc ggc gag tat ggt ttc ccc ctc aag tta aaa gcc cgc tat gac 99
 Lys Pro Gly Glu Tyr Gly Phe Pro Leu Lys Leu Lys Ala Arg Tyr Asp
 15 20 25

15 aac ttt att ggc ggc gaa tgg gta gcc cct gcc gac ggc gag tat tac 147
 Asn Phe Ile Gly Gly Glu Trp Val Ala Pro Ala Asp Gly Glu Tyr Tyr
 30 35 40

cag aat ctg acg ccg gtg acc ggg cag ctg ctg tgc gaa gtg gcg tct 195
 Gln Asn Leu Thr Pro Val Thr Gly Gln Leu Leu Cys Glu Val Ala Ser
 20 45 50 55

tcg ggc aaa cga gac atc gat ctg gcg ctg gat gct gcg cac aaa gtg 243
 Ser Gly Lys Arg Asp Ile Asp Leu Ala Leu Asp Ala Ala His Lys Val
 60 65 70

aaa gat aaa tgg gcg cac acc tcg gtg cag gat cgt gcg gcg att ctg 291
 25 Lys Asp Lys Trp Ala His Thr Ser Val Gln Asp Arg Ala Ala Ile Leu
 75 80 85 90

ttt aag att gcc gat cga atg gaa caa aac ctc gag ctg tta gcg aca 339
 Phe Lys Ile Ala Asp Arg Met Glu Gln Asn Leu Glu Leu Leu Ala Thr
 95 100 105

	gct gaa acc tgg gat aac ggc aaa ccc att cgc gaa acc agt gct gcg	387
	Ala Glu Thr Trp Asp Asn Gly Lys Pro Ile Arg Glu Thr Ser Ala Ala	
	110 115 120	
	gat gta ccg ctg gcg att gac cat ttc cgc tat ttc gcc tgg tgt att	435
5	Asp Val Pro Leu Ala Ile Asp His Phe Arg Tyr Phe Ala Ser Cys Ile	
	125 130 135	
	cgg gcg cag gaa ggt ggg atc agt gaa gtt gat agc gaa acc gtg gcc	483
	Arg Ala Gln Glu Gly Gly Ile Ser Glu Val Asp Ser Glu Thr Val Ala	
	140 145 150	
10	tat cat ttc cat gaa ccg tta ggc gtg gtg ggg cag att atc ccg tgg	531
	Tyr His Phe His Glu Pro Leu Gly Val Val Gly Gln Ile Ile Pro Trp	
	155 160 165 170	
	aac ttc ccg ctg ctg atg gcg agc tgg aaa atg gct ccc gcg ctg gcg	579
	Asn Phe Pro Leu Leu Met Ala Ser Trp Lys Met Ala Pro Ala Leu Ala	
15	175 180 185	
	gcg ggc aac tgt gtg gtg ctg aaa ccc gca cgt ctt acc ccg ctt tct	627
	Ala Gly Asn Cys Val Val Leu Lys Pro Ala Arg Leu Thr Pro Leu Ser	
	190 195 200	
	gta ctg ctg cta atg gaa att gtc ggt gat tta ctg ccg ccg ggc gtg	675
20	Val Leu Leu Leu Met Glu Ile Val Gly Asp Leu Leu Pro Pro Gly Val	
	205 210 215	
	gtg aac gtg gtc aat ggc gca ggt ggg gta att ggc gaa tat ctg gcg	723
	Val Asn Val Val Asn Gly Ala Gly Gly Val Ile Gly Glu Tyr Leu Ala	
	220 225 230	
25	acc tgg aaa cgc atc gcc aaa gtg gcg ttt acc ggc tca acg gaa gtg	771
	Thr Ser Lys Arg Ile Ala Lys Val Ala Phe Thr Gly Ser Thr Glu Val	
	235 240 245 250	

	ggc caa caa att atg caa tac gca acg caa aac att att ccg gtg acg	819
	Gly Gln Gln Ile Met Gln Tyr Ala Thr Gln Asn Ile Ile Pro Val Thr	
	255 260 265	
	ctg gag ttg ggc ggt aag tcg cca aat atc gtc ttt gct gat gtg atg	867
5	Leu Glu Leu Gly Gly Lys Ser Pro Asn Ile Val Phe Ala Asp Val Met	
	270 275 280	
	gat gaa gaa gat gcc ttt ttc gat aaa gcg ctg gaa ggc ttt gca ctg	915
	Asp Glu Glu Asp Ala Phe Phe Asp Lys Ala Leu Glu Gly Phe Ala Leu	
	285 290 295	
10	ttt gcc ttt aac cag ggc gaa gtt tgc acc tgt ccg agt cgt gct tta	963
	Phe Ala Phe Asn Gln Gly Glu Val Cys Thr Cys Pro Ser Arg Ala Leu	
	300 305 310	
	gtg cag gaa tct atc tac gaa cgc ttt atg gaa cgc gcc atc cgc cgt	1011
	Val Gln Glu Ser Ile Tyr Glu Arg Phe Met Glu Arg Ala Ile Arg Arg	
15	315 320 325 330	
	gtc gaa agc att cgt agc ggt aac ccg ctc gac agc gtg acg caa atg	1059
	Val Glu Ser Ile Arg Ser Gly Asn Pro Leu Asp Ser Val Thr Gln Met	
	335 340 345	
	ggc gcg cag gtt tct cac ggg caa ctg gaa acc atc ctc aac tac att	1107
20	Gly Ala Gln Val Ser His Gly Gln Leu Glu Thr Ile Leu Asn Tyr Ile	
	350 355 360	
	gat atc ggt aaa aaa gag ggc gct gac gtg ctc aca ggc ggg cgg cgc	1155
	Asp Ile Gly Lys Lys Glu Gly Ala Asp Val Leu Thr Gly Gly Arg Arg	
	365 370 375	
25	aag ctg ctg gaa ggt gaa ctg aaa gac ggc tac tac ctc gaa ccg acg	1203
	Lys Leu Leu Glu Gly Glu Leu Lys Asp Gly Tyr Tyr Leu Glu Pro Thr	
	380 385 390	

	att ctg ttt ggt cag aac aat atg cgg gtg ttc cag gag gag att ttt	1251
	Ile Leu Phe Gly Gln Asn Asn Met Arg Val Phe Gln Glu Glu Ile Phe	
	395 400 405 410	
	ggc cgg gtg ctg gcg gtg acc acc ttc aaa acg atg gaa gaa gcg ctg	1299
5	Gly Pro Val Leu Ala Val Thr Thr Phe Lys Thr Met Glu Glu Ala Leu	
	415 420 425	
	gag ctg gcg aac gat acg caa tat ggc ctg ggc gcg ggc gtc tgg agc	1347
	Glu Leu Ala Asn Asp Thr Gln Tyr Gly Leu Gly Ala Gly Val Trp Ser	
	430 435 440	
10	cgc aac ggt aat ctg gcc tat aag atg ggg cgc ggc ata cag gct ggg	1395
	Arg Asn Gly Asn Leu Ala Tyr Lys Met Gly Arg Gly Ile Gln Ala Gly	
	445 450 455	
	cgc gtg tgg acc aac tgt tat cac gct tac cgg gca cat gcg gcg ttt	1443
	Arg Val Trp Thr Asn Cys Tyr His Ala Tyr Pro Ala His Ala Ala Phe	
15	460 465 470	
	ggt ggc tac aaa caa tca ggt atc ggt cgc gaa acc cac aag atg atg	1491
	Gly Gly Tyr Lys Gln Ser Gly Ile Gly Arg Glu Thr His Lys Met Met	
	475 480 485 490	
	ctg gag cat tac cag caa acc aag tgc ctg ctg gtg agc tac tcg gat	1539
20	Leu Glu His Tyr Gln Gln Thr Lys Cys Leu Leu Val Ser Tyr Ser Asp	
	495 500 505	
	aaa ccg ttg ggg ctg ttc taagagctcg aattcgc	1574
	Lys Pro Leu Gly Leu Phe	
	510	

<210> 8
 <211> 512
 <212> PRT
 <213> Escherichia coli

5 <400> 8

Met Thr Asn Asn Pro Pro Ser Ala Gln Ile Lys Pro Gly Glu Tyr Gly
 1 5 10 15

Phe Pro Leu Lys Leu Lys Ala Arg Tyr Asp Asn Phe Ile Gly Gly Glu
 20 25 30

10 Trp Val Ala Pro Ala Asp Gly Glu Tyr Tyr Gln Asn Leu Thr Pro Val
 35 40 45

Thr Gly Gln Leu Leu Cys Glu Val Ala Ser Ser Gly Lys Arg Asp Ile
 50 55 60

Asp Leu Ala Leu Asp Ala Ala His Lys Val Lys Asp Lys Trp Ala His
 15 65 70 75 80

Thr Ser Val Gln Asp Arg Ala Ala Ile Leu Phe Lys Ile Ala Asp Arg
 85 90 95

Met Glu Gln Asn Leu Glu Leu Leu Ala Thr Ala Glu Thr Trp Asp Asn
 100 105 110

20 Gly Lys Pro Ile Arg Glu Thr Ser Ala Ala Asp Val Pro Leu Ala Ile
 115 120 125

Asp His Phe Arg Tyr Phe Ala Ser Cys Ile Arg Ala Gln Glu Gly Gly
 130 135 140

Ile Ser Glu Val Asp Ser Glu Thr Val Ala Tyr His Phe His Glu Pro
 25 145 150 155 160

	Leu Gly Val Val Gly Gln Ile Ile Pro Trp Asn Phe Pro Leu Leu Met		
	165	170	175
	Ala Ser Trp Lys Met Ala Pro Ala Leu Ala Ala Gly Asn Cys Val Val		
	180	185	190
5	Leu Lys Pro Ala Arg Leu Thr Pro Leu Ser Val Leu Leu Leu Met Glu		
	195	200	205
	Ile Val Gly Asp Leu Leu Pro Pro Gly Val Val Asn Val Val Asn Gly		
	210	215	220
	Ala Gly Gly Val Ile Gly Glu Tyr Leu Ala Thr Ser Lys Arg Ile Ala		
10	225	230	235 240
	Lys Val Ala Phe Thr Gly Ser Thr Glu Val Gly Gln Gln Ile Met Gln		
	245	250	255
	Tyr Ala Thr Gln Asn Ile Ile Pro Val Thr Leu Glu Leu Gly Gly Lys		
	260	265	270
15	Ser Pro Asn Ile Val Phe Ala Asp Val Met Asp Glu Glu Asp Ala Phe		
	275	280	285
	Phe Asp Lys Ala Leu Glu Gly Phe Ala Leu Phe Ala Phe Asn Gln Gly		
	290	295	300
	Glu Val Cys Thr Cys Pro Ser Arg Ala Leu Val Gln Glu Ser Ile Tyr		
20	305	310	315 320
	Glu Arg Phe Met Glu Arg Ala Ile Arg Arg Val Glu Ser Ile Arg Ser		
	325	330	335
	Gly Asn Pro Leu Asp Ser Val Thr Gln Met Gly Ala Gln Val Ser His		
	340	345	350

Gly	Gln	Leu	Glu	Thr	Ile	Leu	Asn	Tyr	Ile	Asp	Ile	Gly	Lys	Lys	Glu		
		355						360					365				
Gly	Ala	Asp	Val	Leu	Thr	Gly	Gly	Arg	Arg	Lys	Leu	Leu	Glu	Gly	Glu		
		370						375					380				
5	Leu	Lys	Asp	Gly	Tyr	Tyr	Leu	Glu	Pro	Thr	Ile	Leu	Phe	Gly	Gln	Asn	
		385					390					395			400		
Asn	Met	Arg	Val	Phe	Gln	Glu	Glu	Ile	Phe	Gly	Pro	Val	Leu	Ala	Val		
				405					410					415			
10	Thr	Thr	Phe	Lys	Thr	Met	Glu	Glu	Ala	Leu	Glu	Leu	Ala	Asn	Asp	Thr	
				420					425					430			
Gln	Tyr	Gly	Leu	Gly	Ala	Gly	Val	Trp	Ser	Arg	Asn	Gly	Asn	Leu	Ala		
		435					440						445				
Tyr	Lys	Met	Gly	Arg	Gly	Ile	Gln	Ala	Gly	Arg	Val	Trp	Thr	Asn	Cys		
		450				455					460						
15	Tyr	His	Ala	Tyr	Pro	Ala	His	Ala	Ala	Phe	Gly	Gly	Tyr	Lys	Gln	Ser	
		465				470					475			480			
Gly	Ile	Gly	Arg	Glu	Thr	His	Lys	Met	Met	Leu	Glu	His	Tyr	Gln	Gln		
				485					490				495				
20	Thr	Lys	Cys	Leu	Leu	Val	Ser	Tyr	Ser	Asp	Lys	Pro	Leu	Gly	Leu	Phe	
				500					505				510				

<210> 9

<211> 5267

<212> DNA

<213> Klebsiella pneumoniae

<220>

<223> Location complement 300..2153

<220>

<223> Location complement 2166..2591

5 <220>

<223> Location complement 2594..3034

<220>

<223> Location complement 2191..4858

<400> 9

10 agcgctatat gcgttgatgc aatttctatg cgcaccogtt ctocggagcac tgtccgacog 60
 ctttgccgcg cgcacagtcg tgcctcgttc gctacttgga gccactatcg actacgcgat 120
 catggcgacc acaccogtcc tgtggatctc ccactgacca aagctggccc cggcgaccog 180
 cagcgacogc agcgagcccg cgcgaagaa aatgagcaat ccggtgccaa gaaactcgcg 240
 cagcgactgc ccggttaagg tagaagtctg gttcattatc ggcactcctga aatagcacgt 300
 15 taaagagaga ggctggcgcg agcgcccggt taattcgctt gaccggccag tagcagcccg 360
 gtggcgacog cattgcgcgg ccttctctgtt ccccgaaatat tgcctgtccc ggcgaccacg 420
 ccatagtgcg acaaggcttc cgtgataaag tcggggatct caaagtcacg cgatgagcgg 480
 cccaccagca ccacaaaggc gatatcgoga atggaaccgc cgggtgagac ctggcgacgc 540
 gcgogcagcg agttgggtgac aaacacttcc tctttcgctt gcggcgacg gagacgaatt 600
 20 ttttccagcg ggctggcggt atcgatcgcg accagttcgc cctccttgat gtacaccact 660
 ttggcgaaaca ccgcccggct gagggcttcc cgaagaagct ccaccgcgcc attctcgtga 720
 cgaataactga acaggcttcc cactttggcc agcgggtatt tttttatcgc ttccgcccagc 780
 gaaagatcct cgaggcccg ctcgggttta atcaacagcg tgaccatatt ccccgcccg 840
 gcgagatgga ccgcccgttat ctgcccctcc gcgttgacga tcgcccgcac cgtcgagcgg 900
 25 gcgcccagggt cgaggatcgc cagcgcgccc gcacagccgg gagtgggttaa cgccccggcg 960
 atggccatgt tggcctccac gcccccacc accacctcgg tctgcagtcg ggcgtcag 1020
 tcggggcgga taacctgcac ttgcagacga tccgcttcca ccacgcgcg catcccgacg 1080
 gcattctcca tggcgccactc ccgggccatc ccgcccctgca cttgcgcgg aataaacgta 1140
 tccaccgcca gcagatcctg gatgtatata gcgctcatct catggccggt cagggacgcc 1200
 30 attaccttgc gacccgcctc aagcatgcgg ccggcggtgg tgcccggttc gccggcgatg 1260
 tcgcgtacgg gagcgacggc gtcacatgcc tgcatgatgg ctcccgccgc ctggcgacga 1320
 tcggcctctc cgcggcgctt ttccgcgcta atgtagaggt tgcggccggg gatcaccgcg 1380
 gactgcacat cccctcgcg ggtcttgagc accaccgcgg aacgggttgc aatcagggcg 1440

cgggcatggt ggacgatggc ctgggtctct tccgggctta gccgaagaa ggtggcgatc 1500
 ccgtagggat tcgacaggat ccgcaccacc tggcccgccg cggccaattc caccgcggcc 1560
 attaccacct cggggacctg ctccagcagc gtcaactcat ccaccaccgg cagggtttta 1620
 cgcaggcggt tgttcaaccg cagcccgctg tcctttttga ggatcgccgc caccacgttg 1680
 5 atcccccggt cgagcgctc attgagccac cacacggcgt caaggaatc gacggcgctg 1740
 tcaatcagta cgateccacc ctgggcatac tgcgcggccg gcagcgctgc cagcgcggcc 1800
 agggcgatag tgcgtcccac gccaacgccc acccgcggcg gcgtctgcgg gtatgacgg 1860
 atcatggatg attcgggtgat aatgggtctg gtgatggctt ccatacgccac atcgccaatc 1920
 accggcgccg cttcgttaag atagatgcga gagacatgc tcatacgcca cgggtgtttc 1980
 10 gccaggccct gctccagcgc ggcgagggtc ccggcgatat tgtcccgctt ccttttcattg 2040
 ccgctcgctg cgacgatccc gctggcaaca aacgcctctg cctgcgggta gtcggacgcc 2100
 agcgccacct cgggtggggc gttgccgata tcaatcccg ctaataacgg catgctgacc 2160
 tccgcttagc ttcttttacg cagcttatgc cgctgctgat acacttcgcg cgactcccg 2220
 acaaaggcgg cattcactgt cgcagccag gtgtgtcca gctcgctggc gatcgccagc 2280
 15 agctccgctt gcgaggagcg gaacgggcgc agcgcgctt atagatccag aatcgctcg 2340
 tcaggaaatgg cgataagctc cgcgcgcggc cggaaattgc gcgccacgc atggcgctgc 2400
 atctgctcgg caatctgcgc ctgggtactca aggggtctggc gggagatccg cacatcctgc 2460
 gggccacctc cgcagagag caccttctcg agggtaatat cggtaaatg tttgccggta 2520
 ggcgtcagga tatgctccgg gcagcgggtg gctaacggat aatcctgcac gcgcatggtt 2580
 20 ttctcgtcga tggtaactcc cttaactaag cgatgtgcag ggtgacgggc tcgcgctcct 2640
 gcaccacatg tttgtctctt ttgatatgaa atagcgcggc tttggccata aatttcggcc 2700
 gcaccatctg atcgttcacc accggcaccg gcgaagggtg ctctttgcgc gcatagcgcg 2760
 cagcggtttt gccaatctgc cggtaggtct ccagcgctcag cagcgcgccg tgggagaaca 2820
 gctccagggt gctgagcggc agcagatcgc gctgatggat gaccgtgctt ccttcgact 2880
 25 ggataccgat gctgatcccc gagccgctca ggttggccgc atcccgagcc ataaaggaga 2940
 cgtcggaagt gcgcagaatg cgcaccaccg gggcggtgaag cccctctctt tccaccgccg 3000
 caatcagctc tttgaggatc gcgccatggg gcatatcgat cagagtgtga tgcgtggtgtt 3060
 tatcgaaggc agggccgacg ccgatcacca cttcatcggc gcgttcacgc gcagaagcta 3120
 cccgcgcctc gcgggttttc agggtaaaag agggctgaat ttgggtttgc tgttgacag 3180
 30 gaataccgcc ttattcaatg gtgtcgggct gaaccacgcc cggaaatatt ttgatctccg 3240
 cccagcgctc ggcagagatg cgateagccg tgcccgccgc ctgatagta ttgatgtcgt 3300
 tgaccgcact caccacctgc aactgcccgt cgagaatggc cgaggtctgc aggtaatcgc 3360
 cggtagcccg ctggcgacgc atattgagaa tattcgtggc gatatactca aagcgctgc 3420
 ggctcagcgc gccgacaata tcgaggccgg tgatgttgcc cttcatcgc tcttcaccg 3480
 35 cactcagatc ctccaccagc ttacgcggcg gcatctcgtt gctgcgctgc gcgtaggtgg 3540
 cggcctccac ctctcgtcgc gcgattggcg gcagcccccag ctcgcggaaa accgcctgga 3600
 tcgccccgcg cgctttctgg cgaatggcaa tggtttcgcg ctcggtcacc ggaagcaggc 3660

cgccgtcaac catcagggtca cgctgcagga tgttgaatc atcaaaatct tccgcatcga 3720
 agttcgagcc gggaacatg ttgtcgtagt tcggcaccgc gctgtagccg gaaaaataa 3780
 agtcgggtgc cggcagcatc tgcacagggt tgcgcgcggt gcggcgaaata tccgagtggtg 3840
 agaaagtctg gtctgtggcg gacgccactt cgaggtcgag catagaggcg atcaggtttt 3900
 5 ccgcaggcac cgcgcgaatg cccgacggca cagcgcggt catgccgata cagctcaccg 3960
 cgccgttttg cagtccttga accccggcgc ctttagtaat gaagatgcag cgcgatttga 4020
 ggttagagcat cgacttgctc tccgaataga ccatcagcgc ttcggatccg gtgcccggagg 4080
 tgttagcgcat tttaaccgc cgggaggcgt aggcgcgaggc gaggaacgcc tttagaccagc 4140
 gcgtatcatc gccgtcggtg aataccgctt cggtgcggtg gaccacacc gtctcggcgt 4200
 10 agctggttaa gccacgcag cccagctcca gctcgggtgc ctcttcacc gagcactgcg 4260
 tcaacacgcc gggggcgccg cactgcgaac cgaccaacag ccgacgggcy ttaaacggcg 4320
 cgtagcgcgc gataccgacc gtggtctcct gttctgagaa gccgcggatc ccggcctcgg 4380
 cggcgtcagc ggcaatctgc accggattat ctttgagatt ggtgacgtg cactggttgg 4440
 aggggggtccg gcgggcacgc atcttctgca gcgccatcat catctccacc acgttcatct 4500
 15 gcgccatcac ctgcaccgct ttggccggcg tgatggcgggt agtgatggca atgatctctt 4560
 cccgggtgac tggaatatcc accagcatac gggctatttc caccgcctcc aggcgcattg 4620
 ctgctctgt gcgtcaacg ttgactgcgt aatcggcgat aaatcggtcg atcatgtcaa 4680
 actggtcccg gcgtttgcg tccagttcga cgatcagacc gttgtccact tttactgaag 4740
 agaccgggtc aaaggggctg tccatggcga tcagccctc ttcaggccac tcgccaatca 4800
 20 gcccgctcgt attgacgggg cgctgggcca gtactgcaaa tcgttttgat cttttcattg 4860
 ttcatcggct caaaaggta aatccgcaga cggtagcgaa tacgcgggc cagcgtcgtt 4920
 gccgcccggc cattaccggc aatagcggaa ctttaaatga gccagtggtg aaaaaataa 4980
 atttaatttc gtttcaattt ggcacacgaa atctaccgac agtttacta tgaacttta 5040
 ctccggcgcg aaaaaataa aatgtgatcg cccgcaatga tataaatcaa ttaataaaaa 5100
 25 acgcccttaa ttacgttttt cgcagcttat tttaacccta ttgactaaat catggcgggc 5160
 gacaaaaata cgctgacaaa aataaagcaa gccaacgaa tggtaaatgt tttttactat 5220
 cgccccctac tgactattcg cgcagcgtt atcctgggtc gggagaga 5268

<210> 10

<211> 607

30 <212> PRT

<213> Klebsiella pneumoniae

<400> 10

Met Pro Leu Ile Ala Gly Ile Asp Ile Gly Asn Ala Thr Thr Glu Val

1

5

10

15

Ala	Leu	Ala	Ser	Asp	Tyr	Pro	Gln	Ala	Arg	Ala	Phe	Val	Ala	Ser	Gly
			20						25					30	
Ile	Val	Ala	Thr	Thr	Gly	Met	Lys	Gly	Thr	Arg	Asp	Asn	Ile	Ala	Gly
			35					40					45		
5	Thr	Leu	Ala	Ala	Leu	Glu	Gln	Ala	Leu	Ala	Lys	Thr	Pro	Trp	Ser
			50				55						60		Met
Ser	Asp	Val	Ser	Arg	Ile	Tyr	Leu	Asn	Glu	Ala	Ala	Pro	Val	Ile	Gly
			65			70					75				80
Asp	Val	Ala	Met	Glu	Thr	Ile	Thr	Glu	Thr	Ile	Ile	Thr	Glu	Ser	Thr
10					85					90					95
Met	Ile	Gly	His	Asn	Pro	Gln	Thr	Pro	Gly	Gly	Val	Gly	Val	Gly	Val
			100						105				110		
Gly	Thr	Thr	Ile	Ala	Leu	Gly	Arg	Leu	Ala	Thr	Leu	Pro	Ala	Ala	Gln
			115				120						125		
15	Tyr	Ala	Glu	Gly	Trp	Ile	Val	Leu	Ile	Asp	Asp	Ala	Val	Asp	Phe
			130				135					140			Leu
Asp	Ala	Val	Trp	Trp	Leu	Asn	Glu	Ala	Leu	Asp	Arg	Gly	Ile	Asn	Val
			145			150					155				160
Val	Ala	Ala	Ile	Leu	Lys	Lys	Asp	Asp	Gly	Val	Leu	Val	Asn	Asn	Arg
20				165					170						175
Leu	Arg	Lys	Thr	Leu	Pro	Val	Val	Asp	Glu	Val	Thr	Leu	Leu	Glu	Gln
			180						185					190	
Val	Pro	Glu	Gly	Val	Met	Ala	Ala	Val	Glu	Val	Ala	Ala	Pro	Gly	Gln
			195					200						205	

	Val Val Arg Ile Leu Ser Asn Pro Tyr Gly Ile Ala Thr Phe Phe Gly		
	210	215	220
	Leu Ser Pro Glu Glu Thr Gln Ala Ile Val Pro Ile Ala Arg Ala Leu		
	225	230	235 240
5	Ile Gly Asn Arg Ser Ala Val Val Leu Lys Thr Pro Gln Gly Asp Val		
	245	250	255
	Gln Ser Arg Val Ile Pro Ala Gly Asn Leu Tyr Ile Ser Gly Glu Lys		
	260	265	270
	Arg Arg Gly Glu Ala Asp Val Ala Glu Gly Ala Glu Ala Ile Met Gln		
10	275	280	285
	Ala Met Ser Ala Cys Ala Pro Val Arg Asp Ile Arg Gly Glu Pro Gly		
	290	295	300
	Thr His Ala Gly Gly Met Leu Glu Arg Val Arg Lys Val Met Ala Ser		
	305	310	315 320
15	Leu Thr Gly His Glu Met Ser Ala Ile Tyr Ile Gln Asp Leu Leu Ala		
	325	330	335
	Val Asp Thr Phe Ile Pro Arg Lys Val Gln Gly Gly Met Ala Gly Glu		
	340	345	350
	Cys Ala Met Glu Asn Ala Val Gly Met Ala Ala Met Val Lys Ala Asp		
20	355	360	365
	Arg Leu Gln Met Gln Val Ile Ala Arg Glu Leu Ser Ala Arg Leu Gln		
	370	375	380
	Thr Glu Val Val Val Gly Gly Val Glu Ala Asn Met Ala Ile Ala Gly		
	385	390	395 400

	Ala Leu Thr Thr Pro Gly Cys Ala Ala Pro Leu Ala Ile Leu Asp Leu		
	405	410	415
	Gly Ala Gly Ser Thr Asp Ala Ala Ile Val Asn Ala Glu Gly Gln Ile		
	420	425	430
5	Thr Ala Val His Leu Ala Gly Ala Gly Asn Met Val Ser Leu Leu Ile		
	435	440	445
	Lys Thr Glu Leu Gly Leu Glu Asp Leu Ser Leu Ala Glu Ala Ile Lys		
	450	455	460
	Lys Tyr Pro Leu Ala Lys Val Glu Ser Leu Phe Ser Ile Arg His Glu		
10	465	470	475 480
	Asn Gly Ala Val Glu Phe Phe Arg Glu Ala Leu Ser Pro Ala Val Phe		
	485	490	495
	Ala Lys Val Val Tyr Ile Lys Glu Gly Glu Leu Val Pro Ile Asp Asn		
	500	505	510
15	Ala Ser Pro Leu Glu Lys Ile Arg Leu Val Arg Arg Gln Ala Lys Glu		
	515	520	525
	Lys Val Phe Val Thr Asn Cys Leu Arg Ala Leu Arg Gln Val Ser Pro		
	530	535	540
	Gly Gly Ser Ile Arg Asp Ile Ala Phe Val Val Leu Val Gly Gly Ser		
20	545	550	555 560
	Ser Leu Asp Phe Glu Ile Pro Gln Leu Ile Thr Glu Ala Leu Ser His		
	565	570	575
	Tyr Gly Val Val Ala Gly Gln Gly Asn Ile Arg Gly Thr Glu Gly Pro		
	580	585	590

Arg Asn Ala Val Ala Thr Gly Leu Leu Leu Ala Gly Gln Ala Asn
 595 600 605

<210> 11

<211> 141

5 <212> PRT

<213> Klebsiella pneumoniae

<400> 11

Met Ser Glu Lys Thr Met Arg Val Gln Asp Tyr Pro Leu Ala Thr Arg
 1 5 10 15

10 Cys Pro Glu His Ile Leu Thr Pro Thr Gly Lys Pro Leu Thr Asp Ile
 20 25 30

Thr Leu Glu Lys Val Leu Ser Gly Glu Val Gly Pro Gln Asp Val Arg
 35 40 45

Ile Ser Arg Gln Thr Leu Glu Tyr Gln Ala Gln Ile Ala Glu Gln Met
 15 50 55 60

Gln Arg His Ala Val Ala Arg Asn Phe Arg Arg Ala Ala Glu Leu Ile
 65 70 75 80

Ala Ile Pro Asp Glu Arg Ile Leu Ala Ile Tyr Asn Ala Leu Arg Pro
 85 90 95

20 Phe Arg Ser Ser Gln Ala Glu Leu Leu Ala Ile Ala Asp Glu Leu Glu
 100 105 110

His Thr Trp His Ala Thr Val Asn Ala Ala Phe Val Arg Glu Ser Ala
 115 120 125

Glu Val Tyr Gln Gln Arg His Lys Leu Arg Lys Gly Ser
 25 130 135 140

<210> 12

<211> 146

<212> PRT

<213> *Klebsiella pneumoniae*

5 <400> 12

Met Pro His Gly Ala Ile Leu Lys Glu Leu Ile Ala Gly Val Glu Glu

1

5

10

15

Glu Gly Leu His Ala Arg Val Val Arg Ile Leu Arg Thr Ser Asp Val

20

25

30

10 Ser Phe Met Ala Trp Asp Ala Ala Asn Leu Ser Gly Ser Gly Ile Gly

35

40

45

Ile Gly Ile Gln Ser Lys Gly Thr Thr Val Ile His Gln Arg Asp Leu

50

55

60

Leu Pro Leu Ser Asn Leu Glu Leu Phe Ser Gln Ala Pro Leu Leu Thr

15 65

70

75

80

Leu Glu Thr Tyr Arg Gln Ile Gly Lys Asn Ala Ala Arg Tyr Ala Arg

85

90

95

Lys Glu Ser Pro Ser Pro Val Pro Val Val Asn Asp Gln Met Val Arg

100

105

110

20 Pro Lys Phe Met Ala Lys Ala Ala Leu Phe His Ile Lys Glu Thr Lys

115

120

125

His Val Val Gln Asp Ala Glu Pro Val Thr Leu His Ile Asp Leu Val

130

135

140

Arg Glu

25 145

<210> 13

<211> 555

<212> PRT

<213> Klebsiella pneumoniae

5 <400> 13

Met Lys Arg Ser Lys Arg Phe Ala Val Leu Ala Gln Arg Pro Val Asn

1

5

10

15

Gln Asp Gly Leu Ile Gly Glu Trp Pro Glu Glu Gly Leu Ile Ala Met

20

25

30

10 Asp Ser Pro Phe Asp Pro Val Ser Ser Val Lys Val Asp Asn Gly Leu

35

40

45

Ile Val Glu Leu Asp Gly Lys Arg Arg Asp Gln Phe Asp Met Ile Asp

50

55

60

Arg Phe Ile Ala Asp Tyr Ala Ile Asn Val Glu Arg Thr Glu Gln Ala

15 65

70

75

80

Met Arg Leu Glu Ala Val Glu Ile Ala Arg Met Leu Val Asp Ile His

85

90

95

Val Ser Arg Glu Glu Ile Ile Ala Ile Thr Thr Ala Ile Thr Pro Ala

100

105

110

20 Lys Ala Val Glu Val Met Ala Gln Met Asn Val Val Glu Met Met Met

115

120

125

Ala Leu Gln Lys Met Arg Ala Arg Arg Thr Pro Ser Asn Gln Cys His

130

135

140

Val Thr Asn Leu Lys Asp Asn Pro Val Gln Ile Ala Ala Asp Ala Ala

25 145

150

155

160

	Glu	Ala	Gly	Ile	Arg	Gly	Phe	Ser	Glu	Gln	Glu	Thr	Thr	Val	Gly	Ile	
					165					170						175	
	Ala	Arg	Tyr	Ala	Pro	Phe	Asn	Ala	Leu	Ala	Leu	Leu	Val	Gly	Ser	Gln	
				180					185						190		
5	Cys	Gly	Arg	Pro	Gly	Val	Leu	Thr	Gln	Cys	Ser	Val	Glu	Glu	Ala	Thr	
			195					200					205				
	Glu	Leu	Glu	Leu	Gly	Met	Arg	Gly	Leu	Thr	Ser	Tyr	Ala	Glu	Thr	Val	
		210					215					220					
	Ser	Val	Tyr	Gly	Thr	Glu	Ala	Val	Phe	Thr	Asp	Gly	Asp	Asp	Thr	Pro	
10	225					230					235					240	
	Trp	Ser	Lys	Ala	Phe	Leu	Ala	Ser	Ala	Tyr	Ala	Ser	Arg	Gly	Leu	Lys	
				245						250					255		
	Met	Arg	Tyr	Thr	Ser	Gly	Thr	Gly	Ser	Glu	Ala	Leu	Met	Gly	Tyr	Ser	
			260					265						270			
15	Glu	Ser	Lys	Ser	Met	Leu	Tyr	Leu	Glu	Ser	Arg	Cys	Ile	Phe	Ile	Thr	
		275						280					285				
	Lys	Gly	Ala	Gly	Val	Gln	Gly	Leu	Gln	Asn	Gly	Ala	Val	Ser	Cys	Ile	
		290				295					300						
	Gly	Met	Thr	Gly	Ala	Val	Pro	Ser	Gly	Ile	Arg	Ala	Val	Leu	Ala	Glu	
20	305					310					315					320	
	Asn	Leu	Ile	Ala	Ser	Met	Leu	Asp	Leu	Glu	Val	Ala	Ser	Ala	Asn	Asp	
				325					330						335		
	Gln	Thr	Phe	Ser	His	Ser	Asp	Ile	Arg	Arg	Thr	Ala	Arg	Thr	Leu	Met	
				340					345						350		

Gln Met Leu Pro Gly Thr Asp Phe Ile Phe Ser Gly Tyr Ser Ala Val
 355 360 365

Pro Asn Tyr Asp Asn Met Phe Ala Gly Ser Asn Phe Asp Ala Glu Asp
 370 375 380

5 Phe Asp Asp Tyr Asn Ile Leu Gln Arg Asp Leu Met Val Asp Gly Gly
 385 390 395 400

Leu Arg Pro Val Thr Glu Ala Glu Thr Ile Ala Ile Arg Gln Lys Ala
 405 410 415

Ala Arg Ala Ile Gln Ala Val Phe Arg Glu Leu Gly Leu Pro Pro Ile
 10 420 425 430

Ala Asp Glu Glu Val Glu Ala Ala Thr Tyr Ala His Gly Ser Asn Glu
 435 440 445

Met Pro Pro Arg Asn Val Val Glu Asp Leu Ser Ala Val Glu Glu Met
 450 455 460

15 Met Lys Arg Asn Ile Thr Gly Leu Asp Ile Val Gly Ala Leu Ser Arg
 465 470 475 480

Ser Gly Phe Glu Asp Ile Ala Ser Asn Ile Leu Asn Met Leu Arg Gln
 485 490 495

Arg Val Thr Gly Asp Tyr Leu Gln Thr Ser Ala Ile Leu Asp Arg Gln
 20 500 505 510

Phe Glu Val Val Ser Ala Val Asn Asp Ile Asn Asp Tyr Gln Gly Pro
 515 520 525

Gly Thr Gly Tyr Arg Ile Ser Ala Glu Arg Trp Ala Glu Ile Lys Asn
 530 535 540

Ile Pro Gly Val Val Gln Pro Asp Thr Ile Glu
545 550 555

<210> 14

<211> 56

5 <212> DNA

<213> Escherichia coli

<400> 14

gctaccatgg cttaacggg accaaggaga tatcatatgt cagtaccogt tcaaca 56

<210> 15

10 <211> 59

<212> DNA

<213> Escherichia coli

<400> 15

gcctcgagtc tagagccgtc gacgggaatt cgagctctta agactgtaaa taaaccacc 59

15 <210> 16

<211> 46

<212> DNA

<213> Saccharomyces cerevisiae

<400> 16

20 gcggtaccaa ggaggtatca tatgttcagt agatctacgc tctgct 46

<210> 17

<211> 33

<212> DNA

<213> Saccharomyces cerevisiae

<400> 17
gogaattcga gctcttactc gtccaatttg gcac 34

<210> 18
<211> 35
5 <212> DNA
<213> Homo sapiens

<400> 18
gcggtaccaa ggaggtatca tatgtcagcc gcgc 35

<210> 19
10 <211> 39
<212> DNA
<213> Homo sapiens

<400> 19
gcgaattcga gctcttatga gttcttctga ggcactttg 39

15 <210> 20
<211> 44
<212> DNA
<213> Escherichia coli

<400> 20
20 gcggtaccaa ggaggtatca tatgaccaat aatccccctt cagc 44

<210> 21
<211> 38
<212> DNA
<213> Escherichia coli

<400> 21
gcgaattcga gctcttagaa cagccccaac ggtttatc 38

<210> 22
<211> 20
5 <212> DNA
<213> Escherichia coli

<400> 22
atcccgccgt taaccacat 20

<210> 23
10 <211> 34
<212> DNA
<213> Escherichia coli

<400> 23
gcggtacat tgttatccgc tcacaattcc acac 34

15 QBMAD\223318

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US00/23878

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : C12P 7/22; C12N 9/02, 9/14, 1/20, 15/00

US CL : 435/155, 189, 195, 252.3, 320.1

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/155, 189, 195, 252.3, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

West; STN files included medline, caplus, uspatfull, embase, scisearch, biosis and biotechds

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	TANG et al. Immunochemical Properties of NAD ⁺ - Linked Glycerol Dehydrogenases from <i>Escherichia coli</i> and <i>Klebsiella pneumoniae</i> . Journal of Bacteriology. December 1982, Vol. 152, No. 3, pages 1169-1174, see the entire article.	1-12
A	BARBIRATO et al. Anaerobic pathways of glycerol dissimilation by <i>Enterobacter agglomerans</i> CNCM 1210: limitations and regulations. Microbiology. 1997, Vol. 143, No. 7, pages 2423-2432.	1-12

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search 16 OCTOBER 2000	Date of mailing of the international search report 20 NOV 2000
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230	Authorized officer TEKCHAND SAJITHA Telephone No. (703) 308-0196